

Who can and cannot be prescribed Super P Force

Is brand Super P Force more effective than the generic in clinical studies?

- [Super P-Force at a Glance: What It Is](#)
- [Quick Facts Digest](#)
- [Steps to Take It](#)
- [Dose guidelines](#)
- [Health Notice of caution and Preventive safeguards](#)
- [Medicines Social exchanges](#)
- [Undesirable effects to look for for](#)
- [Storage infrastructure and Waste Disposal](#)
- [Medicines outside conventional therapy and Similar Therapies](#)
- [Pricing Reference for the United States](#)
- [Regulatory Status and Acquisition in the United States](#)
- [FAQ](#)

Summary: What Super P-Force Is

Super P-Force is a combination oral tablet that pairs two active substances to address two separate male sexual health issues at the same time: erectile dysfunction (ED) and premature ejaculation (PE). A typical formulation contains sildenafil citrate 100 mg, a phosphodiesterase type 5 inhibitor commonly known from Viagra, together with dapoxetine 60 mg, a short-acting selective serotonin reuptake inhibitor used in some countries for PE.

The idea behind the product is to offer erection support from sildenafil and ejaculatory delay from dapoxetine in a single dose. It is important to note that while sildenafil is

approved by the US Food and Drug Administration (FDA) for ED, dapoxetine is not FDA approved. Because of that, Super P-Force in its marketed fixed-dose form is not an FDA-approved product for sale or medical use in the United States. Turn to the section title. [Buying in the USA: Lawful Status and Compliance](#) for more details and guidance about purchasing.

A quick snapshot of the key details

- Standard tablet composition: sildenafil 100 mg plus dapoxetine 60 mg in many marketed versions.
- Active agents: sildenafil citrate (for ED) and dapoxetine (for PE).
- Target conditions: erectile dysfunction and premature ejaculation addressed together.

To jump ahead to practical matters, see [How to Handle It Best](#), our [Dose instructions](#), and safety protocols in [Compliance warnings and Precautionary Measures](#). For legal and purchasing information consult [Legal Eligibility and Purchases in the US](#).

How to take efficiently Super P-Force correctly and safely

General administration recommendations are similar to those for sildenafil monotherapy. Typical instructions include taking one tablet by mouth whole with water approximately 30 to 60 minutes before sexual activity. Avoid crushing or chewing the tablet.

- Timing: Plan dosing about 45 minutes before anticipated activity to allow onset.
- Food: The tablet may be taken with or without food; very fatty meals can slow how quickly sildenafil starts working.
- Hard liquor: Limit alcohol because it can increase dizziness and reduce effectiveness.
- Daily frequency: Do not take more than one tablet within a 24-hour period unless directed by a healthcare professional.

If you are unsure about suitability or the exchange between peoples among teammates fuel progress., provide a full medication list to your clinician before use and consider the results [Important Cautions and Environmental warnings](#) and [Drug Interaction Notifications](#).

Dose instructions and safety limits

The commonly cited strength for many marketed combination tablets is 160 mg total active per tablet (100 mg sildenafil plus 60 mg dapoxetine). Most sources recommend a single tablet as needed and not more than once in any 24-hour period. Always follow a prescriber's directions rather than internet instructions.

If you experience severe or worrying signs after taking the medicine, such as chest pain, fainting, a sudden loss of vision or hearing, severe dizziness, seizures, or an erection lasting longer than four hours, seek emergency medical care. For routine dosing questions, consult your clinician and recheck the [How to Weather It](#) professional guidance.

Notice of risks and preventive steps

- Intended audience: For adult men only. Not approved for women or children under 18.
- Nitrates: Never take with nitrates or nitric oxide donors such as nitroglycerin or isosorbide medications due to risk of severe low blood pressure.
- Other ED treatments: Keep from combining with other PDE5 inhibitors or injectable ED therapies unless your prescriber specifically approves it.
- Cardiac considerations: Persons with significant cardiovascular disease should be evaluated by a clinician before using any ED medication.
- Hepatic and renal impairment: Severe liver or kidney disease may require dose adjustment or avoidance.
- Mental health: Dapoxetine affects serotonin pathways. Monitor for mood changes, agitation, or suicidal thoughts and seek help promptly if they occur.
- Allergies: Don't utilize if you have a known allergy to sildenafil, dapoxetine, or tablet excipients.

For a concise list of interacting medicines, consult [Prescription Drug Contacts besides the quality of the act of interactions matters for trust.](#) For serious adverse events, see [Possible detrimental outcomes.](#)

Drug combination positive communication in actions enhance user satisfaction. to keep in mind

Because Super P-Force combines a PDE5 inhibitor and a serotonergic agent, it has two distinct the act of interaction profiles to watch for.

- Serotonergic drugs: Combining dapoxetine with SSRIs, SNRIs, MAO inhibitors, triptans, tramadol, linezolid, lithium, or St. Johns wort increases the risk of serotonin syndrome.
- CYP3A4 inhibitors: Strong inhibitors such as ritonavir, cobicistat, ketoconazole, itraconazole, and clarithromycin can raise sildenafil and dapoxetine concentrations and may require avoidance or close observation and tracking.
- Alpha-blockers and antihypertensives: Combining these may cause additive drops in blood pressure and dizziness; dose spacing and time sensitive monitoring are important.
- Nitrates and poppers: Organic nitrates or amyl nitrite are contraindicated with PDE5 inhibitors because of potentially life-threatening hypotension.
- Grapefruit products: Grapefruit juice can increase sildenafil levels; discuss avoidance in meeting with your clinician.

Always give your provider a complete list of prescription, over-the-counter, herbal, and recreational substances before starting therapy.

Possible undesirable effects

Common complaints reported with sildenafil and dapoxetine include a dull headache, flushing, nausea, indigestion, dry mouth, back pain, dizziness, fatigue, and drowsiness. Many people tolerate the medicines well, but temporary side effects can occur.

Seek fast medical care for severe signs such as chest pain, sudden loss of vision or hearing, difficulty breathing, fainting, irregular heartbeat, severe swelling, seizures, or an erection lasting more than four hours (priapism). Watch for signs of serotonin syndrome (agitation, confusion, rapid heart rate, sweating, fever, tremors, or muscle stiffness) if taking serotonergic medications.

If symptoms persist or cause concern, contact your healthcare provider. See [Secure data warehouse storage helps organize stock and streamline fulfillment. and Disposal System](#) for safe handling at home.

Temporary physical storage keeps goods safe and organized. and disposal

- Store at room temperature, away from excessive heat, moisture, and direct sunlight.

- Keep tablets in their original blister pack or container until use to minimize exposure to humidity.
- Place all medicines where children and pets cannot reach them..
- Dispose of expired or unwanted medication via community take-back programs or as directed by a pharmacist; do not flush medications down the toilet unless instructed.

Backup options and related medications

Because dapoxetine is not FDA approved, many US clinicians treat ED and PE as separate problems using licensed options for each. Combining approved ED therapy with a PE strategy under supervision is a common approach.

Options for addressing erectile dysfunction

- Tadalafil (Cialis) - longer acting, available for as-needed or low-dose daily use.
- Sildenafil (generic Viagra) - a well-established PDE5 inhibitor widely available in generic form.
- Avanafil (Stendra) - an as-needed PDE5 inhibitor with a relatively quick onset in some users.
- Vardenafil (Levitra) - another PDE5 inhibitor option for as-needed use.
- Nonpharmacologic choices - vacuum erection devices, penile injections under guidance, lifestyle changes, and psychological therapy for performance anxiety.

Premature ejaculation treatment options

- Topical anesthetics - lidocaine or prilocaine formulations applied before intercourse to reduce penile sensitivity.
- Off-label oral antidepressants - some SSRIs such as paroxetine or sertraline are used off-label for PE under clinician oversight.
- Clomipramine - a tricyclic antidepressant sometimes used off-label for PE.
- Behavioral methods - start-stop, squeeze techniques, pelvic floor exercises, and counseling or sex therapy.

When both conditions are present, a prescriber may recommend a licensed PDE5 inhibitor for ED and pair it with an appropriate PE strategy. For approximate costs, see the [Price Reference Guide for the USA](#). For legal factors to evaluate about imported products, see [What is the Legal Position When Buying in the USA](#).

Price guide across the USA (approximate USD)

Prices vary widely by pharmacy, dose, insurance coverage, and discounts. The following are ballpark retail figures to help with comparisons.

- Sildenafil generic (20-100 mg): roughly \$0.50 to \$3.00 per tablet with common discount programs; brand Viagra may cost \$60 to \$80 per tablet without insurance.
- Tadalafil generic (5-20 mg): about \$1.00 to \$4.00 per tablet depending on quantity and discounts; brand Cialis is usually much more expensive.
- Avanafil (Stendra): approximately \$25 to \$70 per tablet depending on dose and drug price list at the pharmacy
- Topical lidocaine-prilocaine products: roughly \$15 to \$40 per small tube or spray bottle.
- Super P-Force via international sellers: often advertised between \$2.00 and \$6.00 per tablet plus shipping, but these sources are unapproved and imports may be legally restricted or risky.

Ask your pharmacist about discount cards, manufacturer coupons, or mail-order options that can reduce out-of-pocket cost. Prices given here are estimates and may change.

The legality and purchasing in the United States

- Prescription requirement: FDA-approved ED medicines such as sildenafil, tadalafil, vardenafil, and avanafil require a valid US prescription. Many states accept telehealth evaluations for a prescription.
- Dapoxetine approval: Dapoxetine is not approved by the FDA. As a result, fixed-dose products that contain dapoxetine plus sildenafil, such as Super P-Force, are not FDA approved for sale in the United States.
- Import rules: The FDA generally prohibits the importation of unapproved prescription medicines. In limited, narrow personal-use exceptions some medications may be allowed, but shipments can be detained or refused and legal risk exists for sellers who import or distribute unapproved drugs.

- Where to purchase: Use licensed US pharmacies or legitimate mail-order pharmacies that require a prescription. Be cautious of websites that do not ask for a prescription or that offer unusually low prices for prescription combinations.
- Safety and quality: Unapproved or foreign-sourced combination products may have variable potency, contamination, or inaccurate labeling. Discuss licensed FDA-approved options and clinically supervised strategies with your healthcare provider.

This page is informational only and does not replace advice from a licensed clinician or legal counsel. For specific legal questions about importation or sale, consult a qualified attorney or the FDA guidance pages.

The questions people ask most

Is Super P-Force sold in US pharmacies?

No. Because dapoxetine is not approved by the FDA, Super P-Force as a fixed-dose combination is not available from licensed US drug retail outlets. See [Similar Medications and Backup Options](#) for approved options.

Can I take Super P-Force if I already take another SSRI?

You should not combine dapoxetine with other serotonergic drugs without medical supervision. The combination can increase the risk of serotonin syndrome. Validating the a description of the intercommunication in [Drug Interaction Notifications](#) and please speak with your prescriber.

What interval is typical may it be taken?

Typically no more than once every 24 hours. Follow your prescriber's instructions and revisit [Dose guidelines](#).

What if I have erectile dysfunction but not premature ejaculation?

An FDA-approved PDE5 inhibitor taken alone, such as sildenafil or tadalafil, may be more appropriate. See choices in [ED treatment options](#).

What should be my next step if I experience secondary consequences?

Contact your healthcare provider for advice about mild symptoms. For emergency signs listed under [Possible outcomes affecting wellbeing and reactions](#), seek immediate medical care.

- [Sartel: Uses, Safety, and Extra Information](#)

- [Can the generic Zoloft match the brand in therapeutic outcomes?](#)
- [Nitrofurantoin: Patient-Centric Guide](#)

[Additional info about Super P Force](#)

© 2026 Kimbrell Veterinary Clinic PC - [Home page](#) - [We're on the map](#)