

Save Big on Super P Force in 2026: Coupons Up to 40% - in the UK

Super P Force: From A to Z Overview

- [Summary with Brief Facts](#)
- [What does Super P-Force hold?](#)
- [Safe Dosing Guidance](#)
- [The Tablet User Guide](#)
- [Possible Adverse Outcomes](#)
- [Drug Interaction Cautions](#)
- [Storage systems and Disposal Operations and Services](#)
- [Guide prices in the United Kingdom](#)
- [Legal and Regulatory governance Observations in the United Kingdom](#)
- [Other Possible Treatments and Similar Options](#)
- [The FAQ and Its Answers](#)

The Macro view and Quick Facts Primer

Below are concise items that summarise this combined product. The tablet brings two active agents together for use by adult men with both erection difficulties and premature ejaculation.

- Typical formulation: one tablet combining sildenafil 100 mg with dapoxetine 60 mg.
- Live units: sildenafil citrate (a PDE5 inhibitor) and dapoxetine (a short-acting selective serotonin reuptake agent).
- Intended condition coverage: erectile dysfunction and premature ejaculation in adult males.

- Not intended for women or for adolescents.

If you are unfamiliar with the medicine, start at the section [What components make up Super P-Force?](#) besides then check [Notes on the United Kingdom regulation and law](#) before considering a purchase.

What elements are part of Super P-Force? and How It Works Explained

This is a fixed-dose combination tablet that aims to address two related sexual health problems at once. It joins two separate modes of action in a single dose.

- Dapoxetine: a rapidly acting serotonin reuptake inhibitor taken on demand to help delay ejaculation and improve perceived control.
- Sildenafil: a phosphodiesterase type 5 inhibitor that assists erectile function by improving penile blood flow when sexual stimulation occurs.

Combining these agents can produce additive benefits, but it can also combine the harmful impacts along with the pattern of interplay between parties determines group dynamics. pose risks of both medicines. Fixed-dose tablets pairing these two ingredients are not licensed for supply in the UK. See [Regulatory and advice related to legality for the UK](#) for details on licensing and lawful supply.

For practical use, continue to [Practical Dosing Guidance](#) and [Tablet Tutorial: How to Get the Most Out of It](#).

Guidelines for Dosing

Typical advertised strength is sildenafil 100 mg plus dapoxetine 60 mg in a single tablet. The common instruction is to take only one tablet within any 24-hour period, as an on-demand dose before sexual activity.

Some people may need a different dose or timing because of age, impaired kidney or liver function, other medical conditions, or concomitant medicines. Older adults and people with organ impairment should seek personalised medical advice. If an experienced an overdose or severe reaction occurs, obtain urgent clinical assistance.

Before using inspect this. the core safety guidelines at [Immediate Drug Interaction Alerts](#) and compare other options at [Options and Comparable Treatments](#).

Getting Started on the Tablet

Practical instructions for on-demand administration:

- Take one tablet about 45 minutes before planned sexual activity, unless a prescriber advises otherwise.
- Swallow the tablet whole using water. Do not crush, split, or chew it.
- This may be taken with or without food. A large, fatty meal can slow absorption and delay onset.
- Limit alcohol intake prior to and during use; alcohol can increase dizziness and reduce effectiveness.
- Avoid grapefruit juice and grapefruit fruit, which can increase sildenafil levels by affecting CYP3A4.
- Effect duration is commonly 4 to 6 hours, though individual response varies.

If you forget a planned dose, do not take more than one tablet in 24 hours. For questions about combining with other therapies, consult [Drug Interaction Alerts Portal and Environmental warnings](#).

Adverse Consequences May Occur

Common, usually transient detrimental outcomes seen with the component medicines include:

- Pain in the head with facial flushing
- Stomach upset, indigestion, or dry mouth
- Giddy feeling, fatigue, or drowsiness
- Lumbar discomfort or nasal congestion

Seek immediate medical attention if any serious or unusual symptoms occur, such as:

- Severe faintness, chest pain, shortness of breath, or irregular heart rhythm
- Sudden loss or change of vision, or sudden hearing loss or persistent ringing
- Recurrent seizures, severe agitation, hallucinations, or marked mood disturbance
- Features suggestive of serotonin syndrome: confusion, high temperature, sweating, shivering, rapid heartbeat, muscle stiffness, or tremor
- Priapism: a painful erection lasting more than four hours

This list does not cover everything. If you have concerns about a symptom, contact your doctor or local emergency services. For safe disposal of unused medicine, see [Storage media and Disposal Operations](#).

Warnings, Conditions precluding use and Social social interactions span conversations, gestures, and feedback. Among Drugs

This product is intended for adult men with documented erectile dysfunction and premature ejaculation. It is not suitable for women, adolescents, or people without these conditions.

- Do not take this tablet on the same day as another erectile dysfunction medicine.
- Avoid if you are taking nitrates or nitric oxide donors, which can cause a dangerous drop in blood pressure when combined with PDE5 inhibitors.
- Keep separate from riociguat or other soluble guanylate cyclase stimulators.
- Inform prescribers about all medicines and supplements: contact and responses across departments improve alignment. are possible with alpha-blockers, potent CYP3A4 inhibitors (some antivirals, macrolide antibiotics, azole antifungals), other antidepressants, and recreational nitrates such as amyl nitrite.
- Concomitant serotonergic medicines raise the risk of serotonin syndrome. Do not use dapoxetine with monoamine oxidase inhibitors or within 14 days of stopping an MAOI.
- Exercise caution if you have a history of seizures, serious cardiac disease, recent stroke or heart attack, significant bleeding disorders, bipolar disorder, or uncontrolled blood pressure.
- If you feel drowsy or dizzy after taking the medicine, do not drive or operate heavy machinery.

See [Potential Negative Effects](#) for more on safety, and survey accordingly [United Kingdom regulatory and law oriented notes and guidance](#) for lawful supply knowledge.

Storage media and Disposal Framework

- Store tablets at normal room temperature away from moisture, heat and direct sunlight.

- Store all medicines in a cabinet that is out of reach and out of sight for children and pets.
- Do not use once the expiry date on the pack has passed. Return unused or expired medications to a pharmacy for safe disposal.

If you are exploring other treatment options juxtapose the options them at [Plan B options and Related Therapeutic Options](#).

UK market indicative price list

Private prices vary by supplier, brand, tablet strength and pack size. The figures below are approximate and in GBP for comparison only.

- Generic sildenafil 50 to 100 mg: around 0.30 to 2.50 per tablet through private pharmacy networks.
- Viagra Connect 50 mg (pharmacy only): typically about 4 to 11 per tablet depending on pack size.
- Generic tadalafil 5 to 20 mg: roughly 0.80 to 3.00 per tablet privately; daily low-dose packs alter per-tablet cost.
- Spedra, Avanafil: commonly 8 to 15 per tablet on the private marketplace.
- Dapoxetine 30 or 60 mg (Priligy or generic where available): often 4 to 12 per tablet from private prescribers.
- Overseas fixed-dose combinations such as Super P-Force: frequently advertised online at about 1.80 to 6.50 per tablet from non-UK suppliers. These products may be unlicensed, illegal to import, or of uncertain quality.

Only purchase prescription medicines from UK-registered pharmacies that require a valid prescription. See [Notes on the United Kingdom regulation and law](#) for more on lawful supply and import rules.

UK Regulatory and Legal Observations

The UK regime for medicines aims to protect patients by requiring appropriate licensing and safe supply channels.

- The Medicines and Healthcare products Regulatory Agency (MHRA) is the regulator for medicines in the UK.

- The General Pharmaceutical Council (GPhC) regulates community and online pharmacies in Great Britain; check a supplier is GPhC-registered and displays contact details.
- Sildenafil is generally prescription-only in the UK, except for the branded 50 mg product sold as a pharmacy medicine after pharmacist assessment.
- Dapoxetine is not sold over the counter; a prescription is needed.
- Fixed-dose products that combine sildenafil and dapoxetine are not licensed in the UK and are not supplied by UK drugstore providers.
- It is unlawful for a UK-based seller to supply prescription medicines without a valid prescription from an appropriate prescriber.
- Bringing prescription-only or unlicensed medicines into the UK from abroad can result in seizure by Border Force; imported products may be counterfeit or substandard.
- If you suspect an adverse reaction to a medicine, report it via the MHRA Yellow Card scheme. In an emergency, call 999.

For safe access, consult your GP or use a regulated online service that employs UK prescribers and dispenses via UK-registered pharmacy chains. For treatment options consult [Another set of options and Equivalent Treatments](#) for price data, see [Approximate price points in the United Kingdom](#).

[Take me back the topmost level](#)

Other Options and Related Treatments

A healthcare professional may recommend one or more of the following options instead of a fixed-dose combination. Many are licensed individually in the UK.

- Tadalafil (generic or Cialis): longer duration, can be used in daily low-dose regimens for steady coverage; onset about 30 minutes.
- Spedra as Avanafil: a rapid-onset PDE5 inhibitor with a potentially shorter window of food intercommunication.
- Vardenafil (Levitra or generic): another PDE5 inhibitor with effects similar to sildenafil for many men.
- Sildenafil alone (Viagra or generic): onset typically 30 to 60 minutes, duration about 4 to 6 hours.

- Dapoxetine alone (Priligy): taken 1 to 3 hours before intercourse to address premature ejaculation.
- Topical therapies for premature ejaculation: lidocaine or lidocaine-prilocaine preparations to reduce penile sensitivity.
- Non-medicine approaches: behavioral techniques, pelvic floor exercises, sex therapy and psychological support can be effective either alone or alongside medicines.
- Combination prescribing: clinicians may prescribe licensed separate medicines together when clinically appropriate rather than an unlicensed fixed combination.

The best option depends on your medical history, other medicines, timing needs, and side-effect tolerance. Speak to a UK prescriber or pharmacist for tailored advice and confirm the results cost comparisons at [Pricing guidelines in the United Kingdom](#).

Questions and Answers for Common Topics

- Can women use Super P-Force? No. This product is intended for adult men only.
- How long are the effects felt? Typical duration reported by many users is about 4 to 6 hours, but this varies between individuals.
- May alcohol be consumed here? Limit alcohol. Excessive drinking can worsen additional effects such as dizziness and reduce desired effects.
- Is it safe to buy online? Only use UK-registered pharmacies that require a valid prescription. Review [UK law and notes on regulatory matters](#).
- What about current antidepressant treatment? Do not start dapoxetine without medical advice because of the networked exchange of ideas and serotonin syndrome risk. See [Drug Interaction Advisory and Heads up](#).

[Navigate back to the highest point](#)

- [Does the generic form of Moduretic yield the same therapeutic benefits as the brand?](#)
- [Before You Take Ampicillin: 7 Adverse effects to Know](#)
- [Eleven Remeron over the counter Additional pathways approved by doctors in 2026](#)

[Additional info about Super P Force](#)