

Cost Per Dose: Extra Super Tadarise - in the UK

Therapeutic Efficacy of Generic Extra Super Tadarise Relative to Other Options

- [Introductory a general overview](#)
- [How It Works](#)
- [Components and Benefits](#)
- [Who is the target user](#)
- [A Guide for Taking](#)
- [Safety measures](#)
- [When this applies, refrain from using it.](#)
- [Possible pharmacologic incidental effects](#)
- [Drug Interaction Risks](#)
- [Dose forgotten](#)
- [An experienced an overdose of medicine](#)
- [Storage area](#)
- [Replacement choices with UK Prices](#)
- [UK Legal Standing and Buying Tips](#)
- [FAQs](#)

Big picture summary

Extra Super Tadarise is a combined oral tablet marketed for adult men who face both erectile dysfunction (ED) and premature ejaculation (PE). It pairs a phosphodiesterase

type 5 inhibitor with a short acting selective serotonin reuptake inhibitor to support erection quality and to help postpone ejaculation during sexual activity.

- Common regimen: taken as needed before planned sexual activity
- Main actives: tadalafil plus dapoxetine
- Geographic focus of this page: United Kingdom (regulations, availability, and approximate promotional pricing)

The information that follows covers the mechanism at work of action, composition, who should be assessed for combined therapy, clinical guidance on dosing, safety measures to factor in, user the dynamics of interactions drive engagement and learning., and purchasing advice. For safety checksthe review cycle the procedures. [Harm prevention measures](#) and [If this condition applies, do not use..](#) [Go go to at the top of the page](#)

How It Operates

The tablet contains two medicines with complementary effects:

- Tadalafil relaxes smooth muscle in the penis by inhibiting PDE5, improving blood flow when sexual stimulation occurs.
- Dapoxetine is a rapidly acting SSRI that increases serotonergic signalling in the brain, which can delay ejaculation and increase control.

Both components require sexual arousal to produce the desired results. Because each drug has its own the process of interaction profileto review the list under [Drug Interaction Prevention](#) before you operate. [Up to at the top](#)

Constituents and Strengths

Typical formulations promoted overseas include:

- Dapoxetine 60 mg tablet combined with tadalafil 40 mg per tablet.

These combination strengths are not licensed pack sizes in the UK. UK prescribers commonly manage the two conditions by prescribing separate licensed products (for example dapoxetine for PE and tadalafil for ED) at appropriate doses. For specifics consult a clinician. [Move to top](#)

Who will benefit from this

This product is intended for adult men, after a clinician has confirmed both of the following:

- Premature ejaculation: ejaculating sooner than desired, often shortly after penetration
- Erectile dysfunction: difficulty obtaining or keeping an erection sufficient for satisfactory sexual activity

A healthcare professional should evaluate cardiovascular status, mental health, and other possible causes before starting a combined regimen. Do not self diagnose or self prescribe. [Jump to top top](#)

How to Take This Step by Step

- Follow the instructions your prescriber has given you exactly. Do not raise the dose without a doctor's advice.
- Timing: usually taken about one hour before sexual activity unless otherwise advised by your clinician.
- Swallow whole with water; do not break, crush, or chew the tablet.
- Food: taking with food is fine, but a large fatty meal may delay the onset of effect.
- Frequency: do not take more than one dose in 24 hours. Tadalafil may remain effective for up to 36 hours for some men, so repeated dosing in a single day is not recommended.
- Beverages with alcohol: avoid heavy drinking as it may increase dizziness and reduce response.
- Grapefruit: avoid large quantities of grapefruit or grapefruit juice due to CYP3A4 the quality of two-way communication in actions matters for trust.. See [Potential Partnered people relating to one anothers are the heartbeat of social platforms. Between Drugs.](#)

If effects are inadequate or possible side effects occur, consult your prescriber rather than altering the dose yourself. [Go back to the very top](#)

Safety protocols

- Adults only: suitable for men aged 18 years and over.

- Blood pressure: the combination can lower blood pressure, particularly when standing up quickly.
- Older patients: men over 65 should discuss risks and benefits; monitoring in real time or dose adjustments may be needed.
- Penile abnormalities: use with caution if you have anatomical penile curvature or Peyronie disease.
- Priapism risk: conditions like sickle cell disease, leukemia, or multiple myeloma increase the chance of prolonged erection.
- Renal or hepatic impairment: significant organ impairment requires specialist input before first use.
- Sensory warning signs: stop treatment and get immediate medical help for sudden vision loss or sudden hearing loss.
- Driving and operating machinery: dapoxetine can cause dizziness, fainting, or drowsiness; check how you react before driving.
- Herbal and over the counter products: tell your clinician about all supplements and herbal remedies to prevent unsafe overlap or interpersonal dialogue and exchanges fuel collaboration and innovation..

Only use after a clinical assessment for ED and/or PE and a medicines review. See [OTC and Prescription Drug Meetings and user the act of connecting with otherss drive engagement and learning.](#) for commonly implicated drugs. [Go to the very top](#)

Refrain from using it when this applies.

- You are allergic to tadalafil, dapoxetine, or any tablet excipient.
- You are younger than 18 years.
- You are taking nitrates for angina or drugs that donate nitric oxide, or guanylate cyclase stimulators like riociguat.
- You have had a recent myocardial infarction, recent stroke, or poorly controlled severe heart disease.
- You have symptomatic low blood pressure or uncontrolled hypotension.
- You have severe liver disease or end stage renal disease unless a specialist advises otherwise.

- You have a current or unstable psychiatric disorder such as uncontrolled bipolar illness.
- You have coagulation problems or an unstable seizure disorder.
- You are taking medicines with strong serotonergic effects that could precipitate serotonin syndrome with dapoxetine; check [Pharmacodynamic Drug Interactions](#).

Not for use alongsideout a formal diagnosis of ED or PE and medical supervision. [Go to the very top](#)

Possible effects to expect

Many users tolerate the combination well. When adverse impact on health occur they are often mild and transient. Call urgent care for chest pain, an erection lasting more than 4 hours (priapism), severe fainting or dizziness, or sudden loss of sight or hearing.

- Cranial pain and facial redness
- Nausea, vomiting, or gastric irritation
- Swirling sensation, drowsiness, or fainting
- Nose congestion with discharge
- Lumbar pain or sore muscles
- Amplified perspiration
- Light headedness due to low blood pressure
- Alterations in sexual desire

This set is not exhaustive. Read the patient leaflet and discuss concerns with the guidance of your prescriber. For life-threatening reactions in the UK, call 999 or attend A&E. [Take me back page top](#)

Prescription Drug Interactions

- Absolute combinations to avoid: nitrates and nitric oxide donors (for example glyceryl trinitrate, isosorbide dinitrate) and riociguat because of severe hypotension risk.
- Serotonergic agents: avoid combining dapoxetine with MAO inhibitors, and be cautious within 14 days of stopping an MAOI. Many antidepressants and other

serotonergic drugs increase the risk of serotonin syndrome (SSRIs, SNRIs, TCAs, mirtazapine, tramadol, triptans, linezolid, lithium, and Hypericum perforatum).

- Antiarrhythmics: certain antiarrhythmics (for example thioridazine or some class IA and class III drugs) can affect cardiac rhythm when combined.
- CNS depressants: benzodiazepines, opioids, some antihistamines, and other sedating agents can increase drowsiness or fainting risk.
- Strong CYP3A4 inhibitors: ketoconazole, itraconazole, clarithromycin, erythromycin, some HIV protease inhibitors, and grapefruit juice can raise tadalafil or dapoxetine blood levels.
- CYP3A4 inducers: rifampicin, carbamazepine, phenytoin, and St Johns wort flower may reduce effectiveness by lowering blood levels.
- Antihypertensives and alpha blockers: additive blood pressure reduction can cause dizziness or fainting; monitor carefully.
- Other ED drugs: do not take multiple PDE5 inhibitors together.

Always provide a full list of current medicines and supplements to the clinician prior to prescription. [Go to top](#)

Missed daily dose

As this product is intended for as needed use, there is no routine dosing schedule. If you forget to take the tablet before sex, take it only if there is still sufficient time for it to work. Do not double the dose to cover a i forgot to take a dose. [Proceed to the next top](#)

Overdose scenario

Excess intake may cause severe unfavorable health effects including prolonged painful erection, marked fall in blood pressure, chest discomfort, palpitations, extreme dizziness, agitation, flushing, dry mouth, nosebleeds, nausea, vomiting, or severe indigestion.

If an a fatal overdose is suspected seek urgent medical attention and bring the medication packaging if available. In the UK call 999 or attend A&E. [Return to the previous page page top](#)

Encrypted physical storage keeps goods safe and organized.

- Store in a closed container at normal room temperature away from moisture.
- Keep away from direct heat and sunlight.
- Keep this product out of reach and out of sight of kids and dogs.

Do not use beyond the expiry date. Return unwanted or expired medicines to a pharmacy for safe disposal. [Go to top](#)

UK Price Information and Options to consider (Indicative)

Prices fluctuate by supplier, brand, strength, and pack size. The figures below are approximate private or retail costs in the UK for quick comparison and are not guaranteed.

- Licensed treatments commonly used:
 - Dapoxetine (Priligy) 30 mg or 60 mg: roughly £3 to £10 per tablet on a private prescription.
 - Tadalafil generic (on demand 10 mg or 20 mg): typically £0.60 to £1.50 per tablet; branded Cialis billing rates can be from around £2 to over £10 per tablet depending on strength and pack.
 - Sildenafil generics (50 mg): around £0.50 to £1.50 per tablet; branded Viagra often sells for roughly £5 to £10 per tablet on private sale.
 - Viagra Connect 50 mg (pharmacy only): commonly about £19 to £25 for 4 tablets.
 - Vardenafil (Levitra) and avanafil (Spedra): typically in the range of £3 to £10 per tablet depending on brand.
- A range of combined products sold overseas (not MHRA licensed):
 - Extra Super Tadarise, Tadapox and similar tadalafil plus dapoxetine tablets: often advertised between about £1.50 and £4.00 per tablet from non UK sellers.
 - Super P Force (sildenafil plus dapoxetine): commonly £1.50 to £4.50 per tablet from overseas vendors.

Because Extra Super Tadarise is not licensed in the UK, clinicians typically prescribe the licensed components separately so that dosing and safety are managed appropriately. Costs on the NHS differ and many of these are prescription only (POM) items. For more on lawful purchasing see [United Kingdom Lawful Condition and Purchasing Guidance](#). [Take me back to top](#)

The legal position in the United Kingdom and Buying Tips

- Prescription status: tadalafil and dapoxetine are prescription only medicines (POM) in the UK. Viagra Connect 50 mg is available as a pharmacy medicine following a pharmacist assessment.
- Licensing: Extra Super Tadarise is not licensed by the MHRA. UK pharmacies normally cannot supply unlicensed combination products unless under specific unlicensed medicine frameworks and with appropriate clinical justification.
- Online buying: use only GPhC registered UK pharmacies when purchasing medicines online. A valid prescription is required for prescription items. Websites selling these drugs without prescription may be operating illegally and could supply counterfeit or unsafe products.
- Importation: parcels of unlicensed prescription medicines from abroad may be intercepted by border authorities. Receipt of such items does not ensure they are authentic or safe.
- Age and suitability checks: pharmacists and prescribers will assess cardiovascular health, mental health history, and realizable drug two-way communication in action alerts before supplying these medicines.
- Sporting considerations: check UKAD or WADA guidance and your sport governing body rules. PDE5 inhibitors are generally permitted but policies can change.

Practical advice: consult your GP, a sexual health clinic, or a registered private prescriber for assessment and to obtain licensed medicines from regulated sources. [Jump to content top](#)

Most Frequently Asked Questions

What duration do the effects have?

Tadalafil may remain effective for up to 36 hours in some men; dapoxetine has a much shorter active window measured in hours. Each person has a unique response time.

Can I drink alcohol when using this tablet?

Moderate alcohol consumption may be acceptable but avoid heavy drinking. Excessive alcohol can worsen dizziness and impair erection response.

Is this product suitable for ladies?

No. The combination is formulated for adult men and is not indicated for women.

Can it be used every day without issue?

This combined product is intended for on demand use. There are licensed daily tadalafil regimens available as single ingredient products; discuss daily therapy options with your clinician if relevant.

For further important safety information see [Drug Interaction Prevention](#). [Go to at the top of the page](#)

- [2026 list of 3 Kamagra Effervescent over the counter Alternatives approved by doctors](#)
- [Unwanted adverse effects, Adverse Reactions, and Other Body Impacts of Doxycycline](#)
- [A Report on P Force Fort: Unwanted reactions and Other Body Effects](#)

[Additional info about Extra Super Tadarise](#)