

Online Availability of Modafresh at Low Price - in Australia

Unpacking Modafresh: can it help you?

- [At a glance](#)
- [Highlights of facts](#)
- [Who is this aimed at](#)
- [How to stomach it](#)
- [Risk reduction measures](#)
- [Who should not make use of it](#)
- [Long term effects](#)
- [Potential potential the pattern of mutual interactions determines group dynamics. among medications](#)
- [Dose left untaken](#)
- [Exceeding a safe dose](#)
- [Storage device](#)
- [Alternative treatments and medicines of a comparable nature](#)
- [Indicative price levels in Australia](#)
- [Lawful status and access in Australia](#)
- [FAQs](#)

High level briefing

Modafresh is a trade name for a product containing modafinil, a prescription medicine that increases wakefulness. It can reduce excessive daytime sleepiness associated with certain disorders but does not cure the underlying condition. Continue primary treatments such as CPAP for obstructive sleep apnoea while using modafinil.

This page summarises practical information on appropriate use, safety precautions, comparable medicines, standard cost range and costs in Australian dollars (AUD), and relevant national rules governing access. Jump straight to [Therapeutic drug and medication the dynamics of interactions at scale reveal systemic patterns.](#), skim the [Short facts](#) you can read about [how modafinil is accessed in Australia](#).

[Return to the position you left top of page](#)

Fast data

- Intended for adults; not licensed for people under 17 years of age based in Australia.
- 100 mg and 200 mg are the standard strengths for tablets.
- Primary effect: promotes alertness in select sleep disorders, not a replacement for sleep.
- Modafinil is the active substance.
- Requires a prescription from an authorised prescriber in the Commonwealth of Australia.

If you are new to modafinil, inspect the sections on [Safety guidelines](#) and [Negative reactions](#) before taking your first dose.

[Scroll to top](#)

Who Modafresh is prescribed for

Health professionals may recommend modafinil for adults who have substantial daytime sleepiness due to one of the following:

- Shift work sleep disorder, designed to support alertness during work shifts.
- Excessive daytime sleepiness due to narcolepsy.
- Persistent daytime sleepiness in obstructive sleep apnoea, once primary treatments (for example CPAP) are in place and optimised.

Modafinil supports symptom control but does not address the root cause. See guidance on dosing. [How to survive it.](#)

[Go to top](#)

What you should know about taking Modafresh

Follow your prescriber precisely. Do not alter dose or timing without medical advice.

- Shift work sleep disorder: typically 200 mg about one hour before the shift begins.
- Narcolepsy syndrome or residual sleepiness from OSA: common starting regimen is 200 mg once in the morning.
- Food: may be taken with or without food; administering it late in the day can interfere with nocturnal sleep.
- Dose modifications: people with significant liver impairment usually require a lower dose; older adults may need slower titration.

Until you know how modafinil affects you, avoid driving and tasks that require full alertness. Review [Safety reminders](#) and [Clinical coadministration online a reciprocal exchanges connect users across distances. between drugss](#) for medicines and conditions that may change dosing or safety.

[Go back to where you left off page top](#)

Prestart safety guidelines

Discuss your full medical history with your prescriber, including any history of:

- Heart disease, chest pain, irregular heartbeat, or poorly controlled high blood pressure
- Psychiatric disorders such as depression, anxiety, psychosis, or bipolar disorder
- Impairment in hepatic or renal health status
- Known substance use disorder or recent heavy alcohol use
- Pregnancy plans, current pregnancy, or breastfeeding

Contraception: modafinil can lower the effectiveness of hormonal contraceptives. Use reliable non-hormonal methods while taking modafinil and for two months after the last dose; discuss options alongside your clinician.

Avoid alcohol while taking modafinil because combined effects may be unpredictable. Stop treatment and seek urgent care if you develop a severe skin reaction, fever with rash, mouth ulcers, swelling of the face or throat, breathing problems, or marked changes in mood or behaviour including suicidal thoughts.

[Up to at the top](#)

Who should not take Modafresh?

- People with a history of a serious allergic reaction to modafinil, armodafinil, or related products (for example, Provigil or Nuvigil).
- Individuals younger than 17 years, unless a specialist advises otherwise.

Do not drive or operate machinery until you understand how the medicine affects you. For legal obligations about driving while taking medicines, see [Legal framework for access in Australia](#).

[Go move to the top tier](#)

Potential reactions to therapy

Many people tolerate modafinil well, but common deleterious health effects include:

- Cranial pain and lightheadedness
- Nausea, upset stomach, diarrhea, or reduced appetite
- Anxiety, nervousness, or agitation
- Difficulty initiating or maintaining sleep
- Nasal stuffiness
- Lower back pain

If you notice chest pain, very fast or irregular heartbeat, severe blistering rash, fever, swelling of the face or tongue, confusion, severe mood disturbance, or suicidal thoughts, seek urgent medical attention. Some negative health impacts may relate to tracking the act of interactions helps measure engagement. with other medicines; see [Drug the act of engaging in dialogue considerations](#).

[Scroll to top](#)

Potential a mutual exchange of ideass are not just about words, but meaning. among prescription medications

Provide a complete medicine list to your prescriber and pharmacist, including over-the-counter drugs and herbal products.

- Induction of CYP3A4 enzymes by modafinil can reduce effectiveness of some drugs, notably hormonal contraceptives, cyclosporine, midazolam and triazolam. Use non-hormonal contraception or discuss a different course of action methods.
- Modafinil may increase levels of medicines metabolised by CYP2C19. Examples include diazepam, propranolol, omeprazole, clomipramine and phenytoin; continuous monitoring or dose changes may be needed.
- Warfarin and other anticoagulants: modafinil can alter anticoagulation control. Check INR more frequently when starting, stopping or changing dose.
- Monoamine oxidase inhibitors (MAOIs): avoid combined use unless managed by a specialist due to potential for adverse people engage through a series of intercommunications..

In moments of doubt, consult your pharmacist. For timing relative to other medicines see [How to stomach it](#) and for contraceptive considerations see [Safety oriented mitigating steps](#).

[Go to top](#)

Untaken dose

If you do not take a dose and it is already late in the day, skip the i forgot to take the dose and resume the normal schedule the next day. Do not take a double dose to make up for a missed one; this increases the risk of insomnia and other lingering effects.

[Up to the first place](#)

Overdose situation

Signs of he overdosed can include agitation, severe insomnia, rapid heart rate, high blood pressure, nausea and restlessness. Dial 000 for emergencies in Australia. For immediate poison advice contact the Poisons Information Centre on 13 11 26.

[Return to the place you were page top](#)

Warehouse seasonal storage spaces provide room for off-season items.

- Keep tablets in their original packaging with the container closed.
- Store at normal room temperature, protected from excessive heat and moisture.
- Ensure that children and pets cannot see or reach this.

If you plan to travel with prescription medicines in review importation rules and carry a copy of your prescription conjoined with the medicine in original packaging. See [Access rights and status of lawful recognition in Australia](#) for more on import documentation.

[Go to the top the page's top](#)

Other options plus analogous medicines

Depending on the diagnosis, individual response and availability, clinicians may consider other options:

- Provigil is the originator brand of modafinil; armodafinil (for example Nuvigil overseas) contains the active enantiomer and is sometimes preferred for longer duration.
- Other wake-promoting agents include solriamfetol and pitolisant; access and approvals in Australia may vary and specialist prescribing is typical.
- Classical stimulant medications such as dexamfetamine or methylphenidate are used under specialist care for narcolepsy in some patients.
- Non-pharmacological measures: sleep hygiene, scheduled naps, light therapy for shift work, and ensuring primary treatment for OSA such as CPAP is optimised.

Ask your sleep specialist or treating doctor which option fits your clinical profile. For a cursory price comparison guide, see [Indicative pricing plan in the Australian market](#).

[Return to the starting point the page top](#)

Price guidance in Australia (AUD)

Retail prices depend on brand, strength, pack size, pharmacy and whether the product is an Australian-registered supply or imported under an approved access route. The following ranges are approximate and provided for general orientation only:

- Modafinil 200 mg tablets, pack of 30: typically around AUD 240 to 400 when supplied as registered brands through Australian pharmaceutical stores.
- Modafinil 100 mg tablets, pack of 30: often between AUD 160 and 300.
- Armodafinil 150 mg tablets, pack of 30: commonly AUD 120 to 300 depending on supplier and brand.
- Overseas-sourced generics or unregistered brands obtained legally under special schemes can vary widely; per-tablet cost may be roughly AUD 2 to 10, plus shipping and any clinical or import fees.

Where the Pharmaceutical Benefits Scheme (PBS) applies for eligible symptoms, patient co-payments are lower but specific authority rules may be required. Always confirm current costs with your local pharmacist and be cautious of unregulated online sellers.

[Return to where you left off the summit of the page](#)

Regulatory status with access rights in Australia

- Scheduling: modafinil is a Schedule 4 (Prescription Only) medicine under the national Poisons Standard.
- Supply: a valid Australian prescription is required to lawfully obtain modafinil from community medicine retailers. Prescribers include specialists and appropriately experienced general practitioners.
- Advertising: prescription medicines may not be promoted to the public. This resource is for education and not promotion.
- Personal importation: under the TGA Personal Import Plan some personal supplies may be imported for individual use (typically up to three months supply) if conditions are met. Keep medicines in original packaging and carry a prescription or supporting documentation for border checks; rules change so verify current TGA guidance before ordering overseas.
- Sport: modafinil may be prohibited in-competition. Athletes should check Sport Integrity Australia and WADA lists and obtain a Therapeutic Use Exemption where appropriate.
- Driving and safety-sensitive work: do not drive or perform safety-critical tasks if you feel drowsy or impaired. Legal responsibility for safe driving remains with the

driver and employers should follow Austroads and state guidance regarding fitness for work.

- Pregnancy and infant feeding: use only when benefits outweigh anticipated risks and with appropriate medical oversight. Effective contraception is recommended during treatment and for two months after stopping if using hormonal contraceptives.

For definitive and up-to-date information consult the Therapeutic Goods Administration (TGA), your state or territory health authority, and your healthcare provider.

[Go back to the number one spot of the page](#)

The most common questions

Does Modafresh stand in for CPAP in obstructive sleep apnoea management?

No. Modafresh may help reduce daytime sleepiness that remains after optimal primary therapy, but it does not treat airway obstruction. Continue CPAP and other prescribed devices unless instructed otherwise by your treating clinician. See [Who this item is for](#).

Is it permissible to drink while on Modafresh?

You should avoid alcohol. Combined use can unpredictably increase adverse outcomes such as drowsiness, impaired judgement or gastrointestinal upset.

How quickly does modafinil work?

Many people report improved alertness within one to two hours after taking a dose. For shift work, dosing about an hour before starting the shift is commonly recommended; see [How to deal with this matter](#).

Is Modafresh suitable for teenagers?

Modafresh is not licensed for people under 17 Australian territory. Any paediatric use should be managed by a specialist with appropriate assessment.

Can I take Modafresh with coffee?

Caffeine may increase agitation, jitteriness or insomnia when combined with modafinil. If you consume caffeine, monitor for amplified stimulant effects and discuss with your prescriber if concerned.

[Go to the very top](#)

Important clarification

This page is intended for general information and does not replace personalised medical advice. Always follow the guidance of your prescriber plus the written information supplied with your medicine. If you have urgent or life-threatening symptoms seek emergency care.

[Return to the main screen the very top](#)

- [Is there a difference in effectiveness between generic and brand Modalert?](#)
- [Is the therapeutic efficacy of generic Ampicillin comparable to the brand?](#)
- [Is Fenbendazole generic as effective as the brand in treating the condition?](#)

[Additional info about Modafresh](#)

© 2026 Kimbrell Veterinary Clinic PC - [Home page](#) - [We're on the map](#)