

# Extra Super Viagra: Online Purchase Direct

## Comprehensive Summary of Extra Super Viagra

- [A Brief Birds eye view and Composition](#)
- [Who is this guide for](#)
- [How to Take It Like a Pro](#)
- [Dosage Amount Guidance](#)
- [Understanding What Active chemical ingredients Are](#)
- [Precautionary safety steps](#)
- [Drug Interaction Risks and Safety](#)
- [Potential harmful consequences](#)
- [Storage systems Care Tips and Guidelines](#)
- [Options, Substitutes, and Comparisons](#)
- [Common USD Prices](#)
- [What is the the legal status in the United States](#)
- [FAQ](#)

## Overall view and Assembly

Extra Super Viagra is marketed as an as-needed oral tablet that pairs two active agents: sildenafil and dapoxetine. The combination is promoted to address erectile dysfunction and symptoms related to premature ejaculation in a single dose. This page summarizes what is known about the product for people in the United States and emphasizes the importance of clinician guidance before operating.

- Administration format: oral tablet taken before sexual activity; sexual stimulation is required for effect.

- Common formulation mentioned online: sildenafil 100 mg combined with dapoxetine 100 mg, though variations exist depending on manufacturer.
- Main intended effects: improving erection quality (sildenafil) and delaying ejaculation (dapoxetine).
- Safety and notes on legality: see the sections on regulations and interacting with others. For US-specific leadership, collaboration with others is the core.

[Return again to this spot top](#)

## Who should use this

The product is intended for adult men who experience erectile dysfunction, and in theory it also targets premature ejaculation due to the dapoxetine component. It is not suitable for people younger than 18 or for women. Because this is a combined therapy, selection should be individualized and made after a medical assessment.

When erectile problems and premature ejaculation coexist, many US clinicians prefer to manage each problem with approved, separate therapies rather than a fixed-dose foreign combination. Take a look at the other selections for approved options and price comparison across products.

[Navigate back to top](#)

## Directions for taking this medicine

Only follow a clinician's instructions. The following are general, non-personalized suggestions commonly provided for as-needed erectile therapies:

- Timing: typically taken about one hour before sexual activity; onset may be as early as 30 minutes and sometimes up to four hours for some people.
- Food: may be used with or without food. A high-fat meal can delay the onset of sildenafil.
- Maximum frequency: do not take more than once in any 24-hour period unless a prescriber gives different directions.
- Missed or unused doses: since it is taken as needed, skip the dose if sexual activity is no longer planned.
- General precautions: swallow with water, limit heavy alcohol consumption, and avoid grapefruit without clinician approval.

For dosing adjustments and special medical conditions, consult the Proposed Dose and Protective measures for safety area sections.

[Go to top](#)

## Dosage and Administration Guidelines

Your prescribing clinician will determine the appropriate strength and dosing regimen. Important principles include:

- Take only the dose prescribed. Do not increase the dose on your own if the response is unsatisfactory.
- Do not use this product together with other PDE5 inhibitors or any other sildenafil-containing product.
- Medical conditions such as liver or kidney impairment, significant heart disease, or blood pressure disorders may require dose modification or a different therapy.
- Because dapoxetine acts on serotonin pathways, mental health history, seizure risk, and other serotonergic medications must be reviewed prior to use.

Report poor effectiveness or adverse symptoms to your prescriber rather than adjusting dose without advice.

[Return to that point the apex of something](#)

## What Makes The active ingredients making up the formulation Tick

This combination contains two distinct medicines with different the machinery behind its:

- Sildenafil: a phosphodiesterase type 5 inhibitor that facilitates erection by increasing blood flow to the penis during sexual stimulation.
- Dapoxetine: a short-acting selective serotonin reuptake inhibitor that has been studied to delay ejaculation; it is not approved by the FDA for this use in the United States.

Because each ingredient has its own the act of interaction profilean in depth review the Drug Interaction Risks section carefully before taking any combination product.

[Move back to the page's top](#)

# Emergency core safety protective protocols

## When Using Is Not Appropriate

- If you are taking any nitrate medications such as nitroglycerin, isosorbide dinitrate, or nitroprusside.
- If you use recreational nitrates (for example, amyl nitrite or butyl nitrite).
- If a clinician has advised you to avoid sexual activity because of cardiovascular disease.
- If you are allergic to sildenafil, dapoxetine, or any ingredient in the tablet.
- If you are already on another PDE5 inhibitor or any product that contains sildenafil.
- If you are taking MAO inhibitors or thioridazine because of dangerous a reciprocal exchanges are the heartbeat of social platforms. with dapoxetine.

## When to Discuss Risks With Your Clinician

- History of prolonged erections or penile abnormalities.
- Common blood disorders include sickle cell disease, leukemia, and multiple myeloma.
- Recent heart attack, stroke, serious arrhythmia, or unstable angina.
- Severe kidney or liver impairment.
- History of epileptic seizures, bipolar disorder, mania, or a tendency to faint.
- Vision or hearing problems, including sudden changes.

Always seek immediate medical attention for severe or alarming symptoms; use the Secondary effects section to recognize emergencies.

[Return to the point you were at the topmost level of the page](#)

## Drug Interaction Notifications

Provide a complete list of prescription drugs, over-the-counter medicines, vitamins, and herbal supplements to your prescriber and pharmacist. Memorable encounters to consider:

- Nitrate medications or nitroprusside: can cause a dangerous drop in blood pressure and must not be combined.
- Alpha-blockers and other antihypertensives: combined effects may lead to symptomatic hypotension.
- Strong CYP3A4 inhibitors such as ketoconazole, itraconazole, clarithromycin, erythromycin, and some HIV protease inhibitors: these may increase sildenafil levels.
- Enzyme inducers like rifampin and certain anticonvulsants: may lower sildenafil exposure and decrease effectiveness.
- Other serotonergic drugs (SSRIs, SNRIs, TCAs, MAOIs, linezolid, lithium, triptans, tramadol): increase the risk of serotonin syndrome when taken with dapoxetine.
- QT-prolonging agents: the combined effect with serotonergic drugs may increase cardiac arrhythmia risk.
- Grapefruit and grapefruit juice: may raise sildenafil concentrations; avoid or follow medical advice.
- Alcoholic beverage and sedatives: may increase dizziness and impair judgment, particularly with dapoxetine.

This list is not comprehensive. Confirm effective interpersonal dialogue and exchanges require clear communication. with a healthcare professional before first use.

[Jump to results the topmost level of the page](#)

## Potential medication negative effects

Some side effects are common and mild, while others are rare and serious. Monitor your symptoms and seek care for worrying signs.

### Common and Usually Mild

- Headache
- Manual flushing or nasal congestion
- Epigastric discomfort or stomach pain

- Vertiginous sensation or lightheadedness
- Nausea, diarrhea, or dry mouth.
- Tiredness or fatigue

### **Serious Reactions Requiring Immediate Help**

- Signs of an allergic reaction: widespread rash, swelling of face or throat, difficulty breathing
- Pain in the chest, syncope, or an irregular heartbeat.
- Sudden vision loss, severe visual changes, or symptoms suggestive of optic nerve damage
- Diminished hearing or ringing in the ears
- Seizure, severe confusion, or weakness on one side of the body
- Prolonged painful erections lasting more than four hours

Report all unexpected or persistent symptoms to your healthcare provider promptly.

[Return to the context the very top](#)

## **File a storage locker can hold tools and supplies. Handling Tips**

- Store at approximately 77 F (25 C) when practical; manufacturers generally accept short excursions outside this temperature.
- Keep the medicine dry and away from heat and direct sunlight.
- Ensure that children and pets cannot see or reach this. Restrooms should not be used for the storage of files is essential for data management..

[Return to the previous step the page top](#)

## **The Field of Different tracks and Comparisons**

In the United States, clinicians usually treat erectile dysfunction and premature ejaculation separately using approved therapies. Accessible different substitutes include:

- Tadalafil (brand name Cialis): known for a longer duration, up to 36 hours; available in both as-needed and daily dosing regimens.
- Sildenafil (brand name Viagra) generic: widely used for ED, typically onset in 30 to 60 minutes with an effect lasting around four hours.
- Vardenafil (Levitra, Staxyn): similar to sildenafil for many patients; formulations vary by pharmacy.
- Avanafil (Stendra): a faster onset for some patients, sometimes effective within 15 to 30 minutes.
- Approaches for premature ejaculation: behavioral techniques, topical anesthetic agents (for example, lidocaine-prilocaine products), and certain SSRIs or TCAs used off-label under medical supervision.

Because fixed-dose sildenafil plus dapoxetine products are not FDA approved, many clinicians will use approved single-agent products or separate prescriptions rather than an unapproved combination tablet.

[Match them up prices before buying side by side across the United States](#) to situate the cost context.

[Go to the very top](#)

## Typical USD Price Range

Estimated out-of-pocket cash prices in the US vary by pharmacy, dose, quantity, and available discounts. Approximate ranges (no insurance) are:

- Generic sildenafil (50 mg to 100 mg): about \$1 to \$10 per tablet; brand Viagra often costs much more, frequently \$70 to \$90 or higher per tablet.
- Generic tadalafil: roughly \$1 to \$8 per 5 mg daily tablet; as-needed 10 mg or 20 mg may be \$2 to \$12 per tablet; brand Cialis typically costs substantially more.
- Vardenafil generics (where available): commonly \$5 to \$15 per tablet; brand price is higher.
- Avanafil (Stendra): often priced between \$30 and \$75 per tablet depending on strength and pharmacy.

Note: Because sildenafil plus dapoxetine combination tablets are not FDA approved in the United States, routine domestic pricing options is not established. International online offers may appear cheaper, but they carry safety and legal risks and may be seized at the border. Discount programs, membership pharmacies, telehealth services, and insurance may dramatically reduce costs.

[Return to where you left off page top](#)

## The status of legality in the United States

- Prescription requirement: FDA-approved PDE5 inhibitors (sildenafil, tadalafil, vardenafil, avanafil) are available only with a valid prescription from a licensed US clinician.
- Approval status: Fixed-dose sildenafil plus dapoxetine tablets are not approved by the FDA for sale in the United States.
- Domestic sale and distribution of unapproved fixed combinations is not permitted, and shipments marketed as such may be seized or refused.
- Import rules: Importing unapproved medications is generally prohibited. In limited circumstances the FDA may exercise enforcement discretion, but that rarely applies when approved domestic variant options are available.
- Telehealth and e-prescribing: lawful in most states when provided by properly licensed clinicians and dispensed through accredited pharmacies; verify credentials and pharmacy licensure.
- Compounding: compounded formulations must comply with state and federal compounding laws and cannot be used to distribute unapproved commercial combinations widely.

Always verify online pharmacies using your state board of pharmacy or the National Association of Boards of Pharmacy resources before purchasing. If in doubt, ask your prescriber or pharmacist for directional guidance.

[Go back to the very top](#)

## Popular Inquiries and Answers

- Is this product FDA approved in the US? No. The fixed-dose combination of sildenafil plus dapoxetine is not FDA approved for sale in the United States. Lay eyes on the portion addressing legal matters for further details.

- Can I use alcohol with this medication? Alcoholic drink should be limited because it can worsen dizziness and fainting risk, especially with dapoxetine. Please consult a professional your clinician.
- What is the length of time the effect persists? Sildenafil typically works for up to about four hours for many people; individual response varies and sexual stimulation is required.
- Is this for women? No. The marketed product targets adult men. It is not intended for use by women or minors.
- What if I already take an SSRI? Combining multiple serotonergic drugs increases the risk of serotonin syndrome. Discuss existing medications with your prescriber before taking any product containing dapoxetine.

#### [Up to the first place](#)

- [Empagliflozin: A Short Guide to What It Is](#)
- [A primer on Hyzaar: what it is and FDA approval?](#)
- [Kamagra Chewable Flavoured: benefits or not for you?](#)

#### [Additional info about Extra Super Viagra](#)