

How to buy Super Cialis in generic form - in Australia

A Closer Look at Super Cialis: Side Effects and Health Impacts

- [Overview and context](#)
- [Legal and Regulatory oversight Notes](#)
- [The Way It Works](#)
- [Dosing and timing](#)
- [How to Absorb It](#)
- [Interactions among users Caused by Drugs](#)
- [Operational Public safety warnings and Cautionary measures](#)
- [Potential safety concerns to be aware of](#)
- [Personal efficient storage helps minimize clutter and maximize space.](#)
- [Other options](#)
- [Prices expressed in AUD](#)
- [Get More Information](#)

Birds eye view

Super Cialis is a combined oral product designed for adult men, intended to address both erectile dysfunction and premature ejaculation with a single tablet. The usual formulation offered by many suppliers pairs tadalafil with dapoxetine in a fixed ratio.

Typical composition seen online and in promotional materials is tadalafil 20 mg together with dapoxetine 60 mg per tablet. The preparation is intended for on-demand use when a prescriber determines a combined approach is appropriate.

Anchored quick links: jump to the [the device that makes it run](#) please audit this. [guidance on dose and timing](#) peer review [central the quality of interactive engagements matters for trust](#). a secondary optionly, read the line [advice for legal decisions for Australia](#).

How this operates

The tablet combines two agents with different targets and time courses.

Tadalafil improves blood flow to the penis by inhibiting phosphodiesterase type 5, which helps achieve and maintain erections when sexual stimulation occurs. Many men experience an effect window that may extend up to about 36 hours.

Dapoxetine is a short acting selective serotonin reuptake inhibitor used as needed to delay ejaculation. It increases serotonergic activity in the central nervous system to help prolong time to climax.

Because onset and duration differ between the two components, coordinate timing carefully. See [how to take pictures](#) and [timing of the dose](#) for practical recommendations.

How to take charge

Only take this medicine if it has been prescribed or recommended by a qualified clinician. General practical guidance for on-demand use follows.

- Swallow tablets whole with water; do not crush or break them unless explicitly instructed by a healthcare professional.
- Plan dosing roughly 30 to 40 minutes before sexual activity. Tadalafil may start to work from around 20 minutes, while dapoxetine often reaches peak effect later, near 40 to 50 minutes.
- A light meal generally will not prevent effect; a large high fat meal can delay absorption.
- Minimise alcohol on dosing days to reduce the risk of lightheadedness and fainting.
- Avoid grapefruit and grapefruit juice because they can interact with the metabolism of one or both drugs.

If you have any uncertainty about timing or food engagement and communications across departments improve alignment. audit the [dosing and administration timing](#) section or consult your prescriber or pharmacist.

Dose and timing

Use the lowest effective dose for the shortest period necessary and follow professional advice tailored to your situation.

- Common fixed-dose product: one tablet containing tadalafil 20 mg and dapoxetine 60 mg taken as needed prior to sexual activity.
- Frequency: do not take more than one tablet in any 24 hour period. Tadalafil persists in the body, so some clinicians may recommend longer spacing between uses for certain patients.
- If you forget and sexual activity remains planned, take a single dose when remembered; do not double the dose.
- People on interacting medicines or with comorbidities may be advised to use separate agents or lower doses instead of the combination.
- Older patients, especially those 65 years and older, may be more sensitive to harmful bodily effects and should be reviewed carefully before using.

Suspect an overdose death? Obtain medical attention urgently. Overdose syndrome symptoms to look for include pronounced dizziness, fainting, or a prolonged and painful erection. See [cautions and risk controls to ensure safety](#) to learn more.

Communication with the addition of the act of interaction

Significant interplay between parties are not just about words, but meaning. can occur. Always review your full medicine list with a prescriber or pharmacist before starting this product.

- Nitrates and nitric oxide donors (for example glyceryl trinitrate, isosorbide compounds, nitroprusside or recreational alkyl nitrites): combining these with tadalafil can cause a dangerous drop in blood pressure and is contraindicated.
- Alcoholic content: may increase the chance of dizziness, fainting, a dull headache and impairs judgment; avoiding or limiting alcohol is prudent.
- Alpha blockers and many antihypertensives: there is an increased risk of low blood pressure; careful keeping track of metrics and timing adjustments may be needed.

- Strong CYP3A4 inhibitors (for example ketoconazole, itraconazole, ritonavir, cobicistat, clarithromycin): these can raise levels of tadalafil and dapoxetine and may require dose changes or avoidance.
- CYP3A4 inducers (for example rifampicin, carbamazepine, phenytoin, Saint John's wort): these may reduce the effectiveness of the drugs.
- Other serotonergic medicines (for example SSRIs, SNRIs, MAO inhibitors, triptans, linezolid, lithium, tramadol): combining with dapoxetine increases the risk of serotonin syndrome.
- Other erectile dysfunction treatments: combining PDE5 inhibitors or other ED therapies is not recommended unless explicitly advised by a doctor.

Advisory on risks and safety measures

Do not use this combination if any of the following apply.

- Known allergy to tadalafil, dapoxetine or any ingredient in the product.
- Severe damage to the liver or kidneys.
- Use of nitrates or recreational nitrites (poppers).
- Recent major cardiac events or unstable cardiovascular conditions, such as a heart attack within the last 90 days, stroke within 6 months, unstable angina, or severe heart failure.

Talk to your clinician before use if you have any of the following conditions or histories.

- Blood disorders, bleeding tendencies, or an active stomach ulcer.
- Penile structural abnormalities or anything that raises the risk of priapism.
- Certain heart conditions, arrhythmias, aortic stenosis, or uncontrolled blood pressure.
- Neurological or psychiatric disorders, a history of seizures or mania, or prior hyponatremia.
- Certain eye disorders or risk factors for non-arteritic anterior ischemic optic neuropathy.

Practical safety points to review:

- Symptoms such as dizziness, sleepiness, blurred vision or sudden weakness are possible; do not drive or operate machinery until you are sure of how the medicine affects you.
- Stop sexual activity and seek urgent medical help if you experience chest pain, severe lightheadedness, fainting, sudden vision or hearing changes, or stroke-like symptoms.
- This medicine does not prevent sexually transmitted infections. Use barrier methods where needed.
- Not permitted for use by women, adolescents, or people under 18.
- If an erection persists beyond 4 hours seek emergency care immediately to avoid lasting damage.

Potential health effects to watch for

Many users do not have deleterious health effects. When they occur they are often mild to moderate and resolve on their own. If symptoms continue or worsen, seek medical advice.

Commonly reported effects include:

- The pain in the head and flushed cheeks.
- Persistent nasal congestion, runny nose.
- Feeling nauseous, a dry mouth, indigestion or diarrhea.
- Giddy feeling, lightheadedness, drowsiness.
- Backache or muscle aches and sleep disturbance.

Serious reactions that require immediate medical attention:

- Prolonged painful erection lasting more than 4 hours (priapism).
- Severe allergic reactions with breathing difficulty, swelling or extensive rash.
- Sudden vision loss or sudden hearing loss, which may be accompanied by ringing in ears and dizziness.

- Signs of serotonin syndrome from dapoxetine when combined with other serotonergic agents: agitation, confusion, fever, sweating, tremor, muscle stiffness and a rapid heartbeat.

Storage area network

Recommended a storage facility serves as a repository for goods awaiting use. careful precautions:

- Keep tablets in their original packaging, protected from moisture and direct sunlight.
- Store at 20 to 25 C (68 to 77 F); avoid excessive heat.
- Keep this item out of the reach and out of sight of both children and pets.

Other avenues

If a fixed dose combined tablet is not suitable, several other options exist for treating erectile dysfunction and premature ejaculation.

Possible avenues for erectile dysfunction treatment

- Tadalafil alone — available as on-demand dosing (10-20 mg) or lower daily doses (2.5-5 mg) for continuous effect.
- Sildenafil (generic or Viagra) — typically shorter duration and widely used as a first line option.
- Vardenafil (Levitra equivalents) — similar onset and effect to sildenafil for some men.
- Avanafil (for example Spedra) — often a faster onset; availability may be limited in some pharmaceutical retailers.
- Non-oral choices — vacuum erection devices, or intracavernosal injections such as alprostadil after specialist review.

Treatment options for premature ejaculation

- Dapoxetine taken on demand where available — short acting SSRI specifically developed for PE in some markets.
- Topical anesthetics — creams or sprays containing lidocaine or lidocaine/prilocaine to reduce sensitivity.

- Behavioural therapies — start-stop, squeeze technique, and pelvic floor exercises can help many men.
- Oral SSRIs used off-label such as sertraline or paroxetine under medical supervision for longer term control.

Choice of therapy should reflect medical history, concomitant medications, side effect profile, and personal preferences. Discuss your options with an Australian registered prescriber or pharmacist.

Prices shown in AUD

Prices vary depending on brand, pack size and whether a product is imported or supplied locally. The figures below are approximate retail ranges in Australian dollars and should be taken as indicative only.

- Generic tadalafil 10-20 mg: roughly AUD 1.50 to 7.00 per tablet in multi-tablet packs; branded Cialis may be about AUD 20 to 30 per tablet in single pack fee schedule.
- Sildenafil 50-100 mg: generic tablets commonly AUD 0.80 to 5.00 each; branded Viagra often AUD 15 to 30 per tablet.
- Spedra, Avanafil 50-200 mg: where available, around AUD 8 to 20 per tablet.
- Topical PE sprays or creams: about AUD 20 to 45 per pack depending on brand and size.
- Dapoxetine 30-60 mg: not routinely supplied in Australia; overseas vendors may list AUD 3 to 10 per tablet but quality and legality may be uncertain.
- Fixed combination products marketed as Super Cialis (tadalafil 20 mg + dapoxetine 60 mg): advertised by some overseas sellers at roughly AUD 6 to 15 per tablet. Availability and legality in Australia differ; see the [legal compliance section](#).

These items are generally not subsidised on the PBS. For exact price and stock status ask a local pharmacist or prescriber.

Australia commentary on legal and regulatory oversight matters

- Tadalafil is classified as a prescription only medicine (Schedule 4) in Australia and requires a prescription from an Australian registered prescriber for lawful supply.

- Dapoxetine and many fixed-dose combinations that include it are not commonly listed on the Australian Register of Therapeutic Goods (ARTG) for routine pharmacy supply.
- Personal importation is possible in limited circumstances under the Therapeutic Goods Administration (TGA) Personal Importation Plan, usually up to a 3 month supply and typically requiring a prescription. Customs, border and quarantine rules also apply and non-ARTG goods may be detained or destroyed.
- Advertising and online sale of prescription medicines is regulated. Overseas internet sellers may offer products of uncertain quality and some items may be counterfeit.
- Telehealth consultations with an Australian prescriber can result in a valid prescription where appropriate, but the product must be supplyable by an Australian pharmacy if it is on the ARTG.

For current and detailed guidance consult the TGA, your state or territory health authority, or a pharmacist before attempting to import or use products not supplied on Australian soil.

More details

- This note relates to erectile dysfunction and premature ejaculation.
- Typical onset and durations: tadalafil effects can start from about 20 minutes and last up to around 36 hours; dapoxetine commonly peaks near 40 to 50 minutes when taken as directed.
- Please refrain from combining. this combination with other ED medicines unless a doctor instructs you to do so.
- If you experience severe reactions seek emergency medical help immediately.

Healthcare professionals can advise whether a combined tablet or separate therapies are the better option for an individual patient. Take another look the [danger alerts](#) and [global teams rely on robust interpersonal engagements](#). sections for safety details.

- [Are generic Cialis Professional together with the branded drug equally potent?](#)
- [Singulair Generic Efficacy in Comparison to Brand-Name Counterparts](#)
- [All About Suhagra: Uses and Common incidental outcomes](#)

[Additional info about Super Cialis](#)