

How to Get Super P Force Online Cheaply - in Australia

Who qualifies for Super P Force therapy

- [Context and a panoramic view of the main points](#)
- [A guide to understanding how it works](#)
- [How to take correctly](#)
- [Dosage](#)
- [Collateral effects](#)
- [Measures for safety and two-way communication in action observation notes](#)
- [Digital cold storage is used for long term preservation of rare items.](#)
- [Other avenues](#)
- [Price levels in Australia](#)
- [The legal status](#)
- [Quick Q and A](#)

Birds eye view: What is Super P-Force?

Super P-Force is a combined oral tablet intended for adult men who have both erectile dysfunction and premature ejaculation. Each tablet typically contains sildenafil citrate 100 mg together with dapoxetine 60 mg, providing an approach that addresses erection support and delayed ejaculation in one dose.

Sildenafil is a phosphodiesterase type 5 inhibitor that helps increase penile blood flow during sexual arousal. Dapoxetine is a short-acting selective serotonin reuptake inhibitor that can extend time to ejaculation and improve control. When used correctly in suitable patients, the combination aims to enhance both erection firmness and ejaculatory latency.

If you are considering purchase or import to Australia double check the regulatory guidance in [Australia's legal position](#) before proceeding.

Facts in brief

- Designed for: adult men with concurrent ED and PE
- Running components: sildenafil 100 mg plus dapoxetine 60 mg
- Typical onset and effect window: often begins within about 45 to 60 minutes; effects commonly last around 4 to 6 hours
- Available strength: fixed-dose 100 mg + 60 mg tablets

Action the structure that supports it

Sildenafil enhances the erectile response by relaxing smooth muscle in penile blood vessels, which permits increased blood inflow in response to sexual stimulation. Dapoxetine works centrally by altering serotonin signaling to prolong ejaculatory latency and improve voluntary control. Combined, they can produce firmer erections and longer duration before climax when appropriate for the patient.

Procedures for taking Super P-Force

Follow the guidance given by your prescriber exactly. The tablet should be swallowed whole with water. It may be taken with or without food; however, a large fatty meal can delay when the medicine begins to work.

- Timing: most men take it about 45 to 60 minutes before sex; some prefer 1 to 3 hours to align dapoxetine peak levels.
- Alcohol content by volume: keep alcohol intake low or avoid it as alcohol can make dizziness or fainting more likely and may worsen erectile performance.
- Alertness: dapoxetine can cause lightheadedness or fainting; avoid driving or using machinery until you know how it affects you.
- Tablet handling: do not break, crush, or chew the tablet; take it intact with a full glass of water.

Turn to the relevant section [Dosage](#) for frequency limits and to [Protective measures](#) for safety screening pre-use.

Dose administration guidelines

The usual regimen is a single tablet containing sildenafil 100 mg and dapoxetine 60 mg taken when required. Take at most one tablet in a 24 hour period.

A treating clinician should assess whether this strength is appropriate for you. Optional solutions include using single-ingredient products or different dosing depending on health conditions, other medicines, and treatment goals.

In case of suspected the person overdosed or severe reactions, seek emergency assistance. If you are in Australia and need urgent help, dial 000 or reach the Poisons Information Centre at 13 11 26.

This product is not intended for daily scheduled dosing. If you miss a planned occasion, do not double the next dose; take it as instructed prior to sexual activity instead.

Safety care steps, contraindication and real time people relating to one anothers enable quick problem solving. carry risks

Super P-Force is intended for adult men only and must not be used by women or those under 18 years of age.

Do not use Super P-Force if you have high blood pressure:

- Are taking nitrates for angina or recreational nitrates (poppers), as severe hypotension can occur.
- Use riociguat or similar soluble guanylate cyclase stimulators.
- I was advised by my clinician to avoid sexual activity due to cardiovascular risk.
- Have recently had a heart attack or stroke, have major heart rhythm problems, or severe liver disease unless cleared by a doctor.
- Plan to combine multiple ED medicines at once without medical supervision.

Take care when using. and medical supervision if you:

- Take alpha-blockers for prostate or blood pressure issues due to risk of symptomatic low blood pressure.

- Are on serotonergic drugs (SSRIs, SNRIs, MAOIs, linezolid, lithium, tramadol, triptans, or St John's wort), because of possible serotonin syndrome when combined with dapoxetine.
- Are taking strong CYP3A4 inhibitors such as ketoconazole, itraconazole, ritonavir, or clarithromycin, which may raise levels of dapoxetine or sildenafil.
- Have a history of fainting, orthostatic hypotension, or certain eye disorders like retinitis pigmentosa.
- Regularly consume grapefruit or grapefruit juice, which can alter the blood levels of these medicines.

Avoid excessive alcohol and very high fat meals around dosing, as these can increase negative effects or delay onset. For rules about supply and importation check [The status of legal recognition in Australia](#).

Possible untoward effects

Many users tolerate this combination without issues; when latent effects appear they are often mild and transient. Standard reactions include:

- Headache discomfort and facial flushing frequently occur
- Feeling off balance or unsteadiness
- Gastric upset, nausea or dry mouth
- Fatigue, drowsiness or back pain

Go to the emergency department for urgent care serious symptoms

- Chest pain, fainting, severe dizziness or difficulty breathing
- Rapid, slow or irregular heartbeat
- Sudden loss of vision or hearing ability., or new tinnitus
- Seizures, extreme mood fluctuations, irritability or suicidal ideation
- Swelling of the extremities
- Priapism: erection lasting more than 4 hours
- Signs of serotonin syndrome: high temperature, sweating, tremor, diarrhea, agitation or confusion

This is merely a partial list. If you are concerned about unwanted undesired reactions, stop the medicine and consult a healthcare professional. Report significant undesired reactions to the Therapeutic Goods Administration (TGA).

Secure data containers and shelves organize storage space effectively. and packaging

Keep tablets in the original packaging at room temperature away from moisture, heat and direct light. In Australia it is commonly advised to store below 25 degrees C unless the label states otherwise. Always store medicines out of reach of children and pets.

Nonconventional medicine and other related remedies

If a fixed-dose combination is not suitable, medical professionals may propose separate agents or different regimens tailored to your condition and interactive engagements among teammates fuel progress..

- Premature ejaculation management options:
 - Dapoxetine as a single-ingredient product (brand example: Priligy where available)
 - Other SSRIs sometimes used off-label under supervision, such as sertraline, paroxetine or fluoxetine
 - Topical anesthetic preparations containing lidocaine or prilocaine
- Treatments for erectile dysfunction:
 - Sildenafil (generic or brand Viagra)
 - Tadalafil (generic or brand Cialis)
 - Vardenafil (Levitra) or avanafil (Spedra) options for erection difficulties
- Other combination strategies:
 - Use separate ED and PE medicines timed appropriately instead of a fixed combo
 - Alternative fixed or mixed combos available overseas, for example tadalafil plus dapoxetine; check Australian regulations and clinical suitability before import

The best choice depends on your health profile what is possible online people relating to one another's connect users across distances., tolerability, convenience and cost. Discuss options with an Australian prescriber to compare benefits and risks.

For a succinct cost comparison, see the illustrated table. [The cost of goods in Australia](#).

Australian pricing approach reference (indicative)

Prices vary widely depending on brand, pharmacy, pack size and whether telehealth or courier fees are involved. The ranges below are approximate and intended for orientation only.

- Generic sildenafil 100 mg at Australian pharmacies: around A\$10 to A\$25 per tablet; bulk packs often reduce the per tablet price.
- Super P-Force 100 mg + 60 mg (commonly imported and not generally stocked by Australian community pharmacies): frequently quoted at about A\$3 to A\$10 per tablet from overseas suppliers, excluding shipping and consultation fees.
- Brand Viagra 100 mg: often about A\$20 to A\$40 or more per tablet depending on pack size and retailer.
- Tadalafil 20 mg generic: roughly A\$5 to A\$15 per tablet; branded Cialis is typically costlier.
- Dapoxetine 60 mg formulation imported: approximately A\$4 to A\$12 per tablet from overseas vendors.

Note: most ED treatments are not subsidised on the Pharmaceutical Benefits Scheme (PBS) for typical indications of, so private payment is common. When buying online, prioritise reputable Australian pharmacies or follow lawful importation routes described in [The standing of the law in Australia](#).

Legal and regulatory considerations in Australia

- Prescription requirements: Sildenafil is classified as a prescription-only medicine (Schedule 4). Dapoxetine is not commonly supplied through Australian pharmacies and may not be registered with the TGA for routine local supply. Prescription medicines should be obtained with a valid Australian prescription.
- Online purchases: Obtaining prescription medicines without a prescription is unlawful. The TGA warns about counterfeit or substandard products sold via unverified websites.
- The Personal Importation Scheme (PIS): Under defined conditions an individual may import up to 3 months supply for personal use. Usually you should have an Australian prescription, keep products in original packaging, not resell them and be prepared to show documentation to Border Force if requested.

- Customs: Shipments may be examined, delayed or seized if they violate PIS conditions, contain unapproved ingredients or pose safety concerns.
- Advertising rules: Public advertising of prescription medicines is restricted within Australia.

Before buying Super P-Force, confirm lawful options with an Australian-registered prescriber or pharmacist and ensure compliance with current TGA and Border Force rules.

Instant FAQ

How long until it takes effect?

Most men feel effects within about one hour. Allowing 1 to 3 hours before sexual activity helps ensure dapoxetine reaches effective levels. A large meal may delay the onset.

Can I take more than one tablet in a day?

No. Do not exceed one tablet in any 24 hour period. If the response is inadequate, speak with your doctor about an alternate path dosing or different medicines. See [Different tracks](#) use this for options.

Who should avoid using Super P-Force?

People taking nitrates or riociguat, those for whom sexual activity is contraindicated, or individuals advised against PDE5 inhibitors or dapoxetine should avoid this medicine. Review full risk mitigation measures at [Precautions plus the act of engaging in dialogues create context for decisions.](#)

[Return the page's top](#)

- [Cialis Professional: Unpleasant Reactions and Additional Body Impacts](#)
- [6 Treatment related secondary effects of Hydrochlorothiazide You Might Experience](#)
- [The Complete Primer on Trazodone](#)

[Additional info about Super P Force](#)