

Viagra Price and Shop Options - in the UK

Viagra: Use Cases, Side Effects, and More

- [Macro view](#)
- [Lawful status within Great Britain](#)
- [What makes it work](#)
- [Administration directions and dosage](#)
- [Drug interpersonal engagement potential](#)
- [Safety care steps](#)
- [Who should refrain from taking it?](#)
- [Expected chronic side effects](#)
- [Dose not administered](#)
- [In an an overdose](#)
- [Storage infrastructure](#)
- [Alternative drugs and similar pharmaceutical options](#)
- [Common UK prices](#)
- [Use in sports](#)

Scope and a survey of the main points

Sildenafil is a medication often known by the brand name Viagra. It belongs to the group of drugs called phosphodiesterase type 5 inhibitors and is commonly prescribed to treat erectile dysfunction in adult men. Different formulations and doses are authorised for pulmonary arterial hypertension, which is managed separately by specialists.

Typical strengths for erectile dysfunction include 25 mg, 50 mg and 100 mg tablets. The active compound is sildenafil, usually presented as sildenafil citrate.

- Main use: treatment of erectile dysfunction in adult males
- Strengths offered for these tablets are 25 mg, 50 mg, and 100 mg.
- Other licensed use: pulmonary arterial hypertension under specialist care
- Formulations: brand-name and generic versions are available

Want details on action, dosing, or risks? See [How the the device works](#), [Directions for dosing](#), and [Drug interpersonal engagement checks](#).

The standing in law in the United Kingdom

In the United Kingdom, most sildenafil products for erectile dysfunction are Prescription Only Medicines (POM). A specific preparation, Viagra Connect 50 mg, is classified as a Pharmacy medicine and may be supplied without a prescription following a pharmacist consultation and assessment of suitability.

- Supply rules: pharmacists will ask about heart health, current medicines, and erectile dysfunction symptoms to ensure safe supply.
- Age limits: Viagra Connect is generally supplied only to those aged 18 or over.
- Online purchases: use only UK-registered providers. Verify GPhC registration and that prescribing follows UK standards. Avoid sites that supply prescription medicines without checks.
- Import rules: small quantities for personal use may be allowed; commercial import needs licences. Border Force may seize unlicensed or unsafe products.
- Safety preventive actions: avoid driving or operating machinery if you experience dizziness or visual disturbances after taking sildenafil.

For cost information, visit the pricing section a nutsell summary of the chapter. [Usual pricing approach levels in the UK](#). For how this process works and how to stomach it, go to [How it operates step by step](#) and [Medication dosing information and administration guidelines](#).

The method of operation

When a man is sexually stimulated, nitric oxide is released in the penis and raises levels of cyclic guanosine monophosphate (cGMP). cGMP relaxes smooth muscle in the

corpus cavernosum, permitting increased blood flow and resulting in an erection. Sildenafil inhibits the enzyme phosphodiesterase type 5 (PDE5) which breaks down cGMP, thereby supporting erection maintenance during sexual stimulation.

Note: sildenafil does not produce an automatic erection; sexual stimulation is required for the medicine to work.

Dosing guidelines and directions

- Typical starting dose for ED: 50 mg taken around 30 to 60 minutes before sexual activity.
- Adjustments to the dose: the dose can be reduced to 25 mg or increased to 100 mg depending on effect and tolerability. Only one dose per 24 hours is allowed.
- Food people relating to one another can be viewed as exchanges of information.: a large, fatty meal can delay onset of action; taking sildenafil on a lighter stomach usually gives a quicker effect.
- Alcoholic drink product: keep alcohol intake moderate. Alcoholic drink product can reduce erectile response and increase indirect effects such as lightheadedness.
- Grapefruit: avoid grapefruit and grapefruit juice because they can raise sildenafil blood levels via the CYP3A4 pathway.

Before taking your dose sildenafil, discuss other medicines and supplements with your pharmacist or prescriber; see [Potential the exchange between peoples are not just about words, but meaning. caused by drugs](#). If you have doubts about sexual activity and underlying heart disease, read [Cautions to observe](#).

Interacting drug effects

Always provide a complete list of medicines, including over-the-counter and herbal products, to the pharmacist or prescriber. Primary the mutual influence during contacts create context for decisions. include:

- Nitrates and nitrite recreational drugs: taking sildenafil with nitrates (for example glyceryl trinitrate or isosorbide mononitrate) or amyl/butyl nitrite can cause a dangerous fall in blood pressure and is contraindicated.
- Strong CYP3A4 inhibitors: medicines such as ketoconazole, itraconazole, ritonavir and certain macrolide antibiotics can increase sildenafil levels; a dose reduction may be needed.

- Alpha blockers: drugs for prostate or blood pressure like doxazosin or tamsulosin can add to blood pressure lowering; careful timing and cautious dosing are advised.
- Other ED drugs: do not fuse these items. sildenafil with tadalafil, vardenafil, or avanafil.
- Antihypertensives and alcohol: combined effects may increase dizziness or fainting risk.

For guidance on food and grapefruit see [Dosing and administration guidelines](#).

Risk mitigation steps

- Hypersensitivity: do not use if you have a known allergy to sildenafil or any tablet ingredient.
- Diseases of the heart and vasculature: sexual activity imposes strain on the heart; seek medical advice if you have unstable angina, recent heart attack or stroke, severe heart failure, uncontrolled high blood pressure, or significant arrhythmias.
- Vision and hearing issues: rare reports exist of sudden vision or hearing loss. Stop the medicine and seek urgent care if either occurs.
- Older adults: they may be more prone to dizziness or blood pressure changes; dose modification might be appropriate.
- The user cohort this is aimed at: ED products are licensed for adult men. Different sildenafil products are used for pulmonary hypertension under specialist supervision.

See details of the absolute exclusions [Who should not consume it?](#). For common adverse events, see [Possible harmful consequences](#).

Who should not take it?

- Anyone currently using nitrates for chest pain or recreational nitrites.
- Patients receiving riociguat (a soluble guanylate cyclase stimulator).
- Individuals instructed to avoid sexual activity because of unstable cardiovascular conditions.
- People with a known allergy to sildenafil or any excipient in the tablet.

- Those already using another phosphodiesterase type 5 inhibitor should not take sildenafil concurrently.

Possible effects to watch for

Hidden effects are usually mild and temporary. Commonly reported reactions include:

- Cephalalgia with red cheeks
- Stuffy nose or nasal congestion
- Abdominal ache or indigestion
- Lightheadedness and dizziness
- Visual disturbances such as altered colour perception or increased sensitivity to light
- Loose stools

Seek immediate medical help if you experience chest pain, fainting, sudden hearing loss, sudden loss of vision, or an erection lasting more than four hours (priapism), which can cause permanent damage.

For how to store items, see the in computing, storage means the capacity to retain data over time. the document section. [Storage room](#). If you fear you have taken more than the recommended amount, read [Overdose](#).

Dose left untaken

Sildenafil for erectile dysfunction is taken as required. In the event that you forget a dose prior to sexual activity, do not take an extra dose later to make up for it. Adhere to the maximum of one dose in 24 hours.

An non-fatal overdose event

If you suspect an overdosed on medication, obtain emergency medical assessment without delay. Symptoms that may indicate overdosed on medication include severe dizziness, fainting, chest pain, a fast or irregular heartbeat, profound nausea, vomiting, or vision problems.

Archive physical storage keeps goods safe and organized.

- Keep away from children and pets and out of their line of sight.
- Store at room temperature, ideally between 20 and 25 degrees C, away from excessive heat and direct sunlight.
- Protect tablets from moisture; do not store in a bathroom.

Options and similar medications

If sildenafil is unsuitable or ineffective, several optional choices exist. Do not merge therapies unless directed by a clinician.

- Tadalafil: available as on-demand doses (for example 10 mg or 20 mg) and also in low-dose daily regimens; tends to have a longer window of action (up to 36 hours).
- Vardenafil: similar onset and duration to sildenafil; common doses include 10 mg and 20 mg.
- Avanafil: may have a faster onset for some users, typical doses range from 50 mg to 200 mg.
- Generic sildenafil: contains the same active ingredient as Viagra and is generally cheaper.
- Non-pharmacological options: vacuum erection devices, psychosexual counselling, and lifestyle changes such as exercise, weight loss, smoking cessation and reducing alcohol.
- Specialist treatments: alprostadil (injection or urethral formulations) or penile prostheses in refractory cases under urology care.

Inspect price comparison guide grid under this specific section [Typical price tags in the United Kingdom](#) and confirm the findings supply routes in [United Kingdom status of lawful recognition](#).

Average cost range options in Britain

Costs vary by brand, pharmacy and pack size. The figures below are approximate private retail british price levels and may change over time. NHS prescription charges differ and availability on the NHS is determined by clinical need and local policy.

- Generic sildenafil (private prescription): roughly 5 to 15 pounds for a pack of 4 tablets, depending on strength and supplier.
- Viagra Connect 50 mg (pharmacy supply): approximately 20 to 25 pounds for 4 tablets, and 35 to 50 pounds for 8 tablets.
- Tadalafil 10 mg or 20 mg (private prescription): around 15 to 30 pounds for 4 tablets.
- Vardenafil 10 mg or 20 mg: often 20 to 35 pounds for 4 tablets.
- Avanafil brand Spedra 50 mg to 200 mg: typically 50 to 70 pounds for 4 tablets.

Note: generic products are usually the most cost effective. In England a prescription fee may apply per item unless you are exempt; prescriptions are free in Scotland, Wales and Northern Ireland. When supplied as a pharmacy-only product like Viagra Connect, the pharmacy retail price usually applies instead of an NHS charge.

For use with sports equipment

Sildenafil is not included on the World Anti-Doping Agency Prohibited List. Athletes should still consult a team doctor before use and verify any sport-specific rules. Evidence that sildenafil enhances athletic performance at sea level is limited; some studies have examined effects at high altitude. Be mindful of ripple effects such as experiencing headaches and dizziness, and avoid unsafe combinations listed in [Potential coadministration possible medication online engagement and communications connect users across distances..](#)

For the full legal and safety statement, see the [Disclaimer of Liability](#) displayed in the page footer.

- [The essentials of Buspar: FDA approval?](#)
- [What Amitriptyline Is Used For, Its Secondary effects, and More](#)
- [Kamagra dose guide: forms, strengths, administration, and more](#)

[Additional info about Viagra](#)