

How To Acquire Super Levitra With Generic Mastercard - in Australia

The Complete Primer on Super Levitra

- [Outline and Composition Framework](#)
- [Who Could Be Taken Into Consideration](#)
- [Safe Use Manual](#)
- [Dosing, Modifications and Communities with special needs](#)
- [Guidelines for Storing](#)
- [Situations that preclude therapy and Safety Protocols](#)
- [Salient Interactions](#)
- [Secondary adverse effects and When to Seek Help](#)
- [Other Options and Comparable Merchandise](#)
- [Prices for Australia \(Estimated\) \(AUD\)](#)
- [Regulatory and import regulations in Australia](#)
- [FAQs](#)

Top level a brief overview of the subject and Composition Framework

Super Levitra is a branded combination tablet marketed in some international online outlets that pairs vardenafil, a phosphodiesterase type 5 inhibitor used to treat erectile dysfunction, with dapoxetine, a short acting selective serotonin reuptake inhibitor commonly prescribed on demand for premature ejaculation in some countries. Typical formulations seen online are 20 mg vardenafil plus 60 mg dapoxetine per tablet.

In Australia, this fixed dose combination is not listed on the Australian Register of Therapeutic Goods (ARTG). The information here is intended to educate and should not replace assessment and advice from an Australian registered prescriber. For specific

primary safety guidelines, see the [Relative Medical reasons to refrain from use and Safety protocols](#) and [Significant Interactions across networks](#) section components.

Beginning time and duration: vardenafil usually has an onset around 30 to 60 minutes and effects may last roughly 4 to 6 hours for many men. Dapoxetine reaches peak blood levels about 1 to 3 hours after dosing and is designed for on demand use rather than daily treatment.

[Return to the starting point page top](#)

Who Might Earn Consideration

Online promotions present Super Levitra as a single tablet addressing both erectile dysfunction and premature ejaculation. In Australia, however, the Therapeutic Goods Administration has not approved this combined product for any indication. Health professionals in Australia may prescribe separate approved treatments when clinically appropriate.

If your main issue is early ejaculation, discuss Australian approved strategies together with your clinician. For erectile dysfunction, see the [Other Choices and Similar Items](#) section for licensed options available locally.

[Take me to on top of the page](#)

How to Access and Use Without Harm

Only take prescription medicines exactly as directed by a clinician. General practical advice commonly given for products combining a PDE5 inhibitor and dapoxetine includes the following points.

- Take one tablet about one hour before planned sexual activity unless your prescriber advises a different timing.
- Swallow the tablet whole and wash it down with water. Do not crush or divide it unless a pharmacist confirms the product is scored and suitable for splitting.
- Do not exceed one dose in any 24 hour period unless your doctor instructs otherwise.
- A high fat meal may delay the onset of the PDE5 inhibitor; an empty stomach can speed absorption for some people.
- Avoid grapefruit and grapefruit juice unless your prescriber confirms they are safe, because they can change drug levels.

- Avoid excessive alcohol before dosing as alcohol can increase dizziness, worsen collateral effects and impair sexual performance.
- If you miss a planned dose and sexual activity is still appropriate, take the dose when remembered but do not take two doses to make up for the missed one.

If you are unsure about timing with other medicines or meals, consult your GP or pharmacist and assess the details this [Vital Interactions](#) the part called section.

[Go to top](#)

Dose Modifications and Adjustments for Groups with special needs

A prescriber may modify dose or dosing interval depending on age, liver or kidney function, and other medications. Typical considerations include:

- Older adults or people with significant liver impairment may need a lower starting dose or longer time between doses.
- Concurrent use with alpha blockers for prostate symptoms or blood pressure control may require careful spacing and a lower initial dose because of the risk of symptomatic low blood pressure.
- Certain strong enzyme inhibitors or interacting drugs may prompt a prescriber to lengthen the minimum interval between doses or avoid the combination.

Never alter dosing without medical advice. If ill effects occur, contact your prescriber to review dose adjustments or nontraditional treatments.

[Up to the uppermost part](#)

Guidelines for Storage infrastructure

- Store tablets at around 25 C. Short excursions from this temperature are generally acceptable; follow the specific product label for exact limits.
- Keep the medicine in its original blister or container to protect from moisture and light until you are ready to use it.
- Protect from excessive heat and keep out of reach of children and pets.

[Return to the earlier section at the top of the page](#)

Cautions against use due to medical reasons and Safety care steps for Use

Do not take this medicine if

- You are using any nitrate medicine such as glyceryl trinitrate or isosorbide dinitrate, or you take recreational amyl nitrite or other poppers.
- You have been advised by a clinician to avoid sexual activity because of serious heart disease.
- You have severe liver impairment, are on dialysis for severe renal failure, have unstable angina, severe heart failure, or uncontrolled blood pressure problems.
- You have congenital long QT syndrome, a known prolonged QT interval, or you take certain antiarrhythmic drugs like amiodarone, quinidine, procainamide, or sotalol.
- You are allergic to vardenafil, dapoxetine, or any ingredient in the tablet.
- You have taken another PDE5 inhibitor recently; do not join these. with sildenafil, tadalafil, avanafil or other vardenafil products.

Talk to your doctor first if you have

- Any other prescription or non prescription medicines, herbal products or supplements, especially serotonergic agents. Glance at the [Foremost Interactions across networks](#) a part of the document.
- Conditions that increase the risk of prolonged erection or priapism, such as sickle cell disease, leukaemia, or multiple myeloma.
- A history of significant eye or ear problems, seizure disorders, bleeding disorders, stomach ulcers, or severe vascular disease.
- A personal or family history of long QT syndrome or other cardiac rhythm disorders.

If any of these issues apply to you, obtain clinical advice before taking a dose. For severe or urgent symptoms, refer to the [Typical indirect effects and When to Get Help](#) a division in the text.

[Go back to where you left off the page's top](#)

Primary Interpersonal engagements

Several medicines and substances can change how vardenafil or dapoxetine work or increase the risk of damaging effects. Discuss all current and recent medicines with your clinician or pharmacist. Critical the exchange that happens when people meets create opportunities for collaboration. include:

- Any nitrates: combining these with a PDE5 inhibitor can produce a dangerous and high potentially life threatening drop in blood pressure.
- Alpha blockers and some antihypertensives: concurrent use can cause symptomatic low blood pressure and fainting.
- Strong CYP3A4 inhibitors such as ketoconazole, itraconazole, certain macrolide antibiotics and some HIV protease inhibitors: these can raise vardenafil concentrations substantially.
- Enzyme inducers such as rifampicin: these may reduce vardenafil exposure and its effectiveness.
- Serotonergic drugs including SSRIs, SNRIs, MAO inhibitors, linezolid, triptans, tramadol and lithium: combining these with dapoxetine can raise the risk of serotonin syndrome and requires medical oversight.
- Certain antiarrhythmics: these may increase the risk of QT prolongation when used with dapoxetine or vardenafil.
- Grapefruit products: these can increase blood levels of some drugs and lead to more minor side effects.

This collection is not exhaustive. Do not start, stop, or change any medicine without professional advice. If you experience symptoms of serotonin syndrome, such as agitation, high temperature, tremor, sweating, diarrhoea or confusion, seek immediate medical attention.

[Go back to the top rank of the page](#)

Adverse effects and When to Get Medical Help

Many people tolerate treatment well, but adverse reactions can occur. Speak with a professional a healthcare professional if you are concerned.

Common negative additional effects

The flush procedure, suffering from a headache, dizziness, nausea, indigestion, nasal congestion, and mild stomach discomfort are among the more frequent effects.

Serious reactions that warrant immediate medical care

If you develop any of the following seek emergency help or go ahead to the nearest emergency department: chest pain, fainting, severe or irregular heartbeat, signs of a severe allergic reaction such as swelling of the face or throat, rash with breathing difficulty, sudden weakness or numbness, sudden loss of vision, sudden hearing loss or severe ringing, seizures, an erection lasting longer than four hours or painful erection, or symptoms suggesting serotonin syndrome (agitation, sudden high fever, severe tremor, persistent vomiting or diarrhoea, confusion).

Until you know how the medicine affects you, avoid driving or operating machinery if you feel dizzy, faint or have vision disturbances. Take a look at the [Regulatory and import requirements in Australia](#) section for driving considerations.

[Take me to the apex](#)

Substitutes and Comparable Products

If a combined product is unavailable or unsuitable, Australian prescribers commonly consider separate licensed therapies and non-drug approaches.

- Tadalafil (brand name Cialis and generics): a PDE5 inhibitor that can be taken as needed or as a low daily dose for ongoing effect.
- Spedra, the brand for Avanafil: a PDE5 inhibitor with a relatively rapid onset for some men.
- Vardenafil (Levitra): the single ingredient found in Super Levitra when prescribed alone.
- Sildenafil (Viagra and generics): widely used on demand PDE5 inhibitor with many generic options.

Non-pharmacological options include vacuum erection devices, intracavernosal injections such as alprostadil, penile implants for refractory cases, lifestyle measures, and psychological or sex therapy. For premature ejaculation alone, prescribers may consider on demand dapoxetine where available, topical anesthetics, or behavioural techniques.

Overseas combination tablets similar to Super Levitra include sildenafil plus dapoxetine (commonly sold as Super P-Force) and tadalafil plus dapoxetine (sold under various trade names). These are generally not ARTG listed and are not routinely supplied by Australian community pharmacy chains.

[Move back to top](#)

Estimated Prices for Australia in AUD

Costs depend on brand, generic status, pack size and supplier. Many erectile dysfunction medicines are prescription-only and not routinely subsidised by the Pharmaceutical Benefits Scheme (PBS) for standard indications, so patients often pay retail prices. Typical retail ranges are approximate and can change:

- Tadalafil 10 to 20 mg (generic): roughly AUD 2 to 8 per tablet in larger packs; small branded packs such as a 4 tablet Cialis pack often cost around AUD 70 to 100.
- Avanafil aka Spedra 100 to 200 mg: four tablet packs commonly retail in the AUD 80 to 110 range.
- Vardenafil (Levitra) 10 to 20 mg: where available, four tablet packs are often about AUD 80 to 100.
- Sildenafil (generic) 50 to 100 mg: bulk discount tablets can be AUD 1 to 5 each; small branded packs like Viagra may cost AUD 60 to 90 for a 4 tablet pack.
- Super Levitra and similar combinations: because these fixed dose combination products are not TGA approved, they are not normally sold by Australian pharmacy networks. Overseas online offers vary widely, for example AUD 1 to 10 per tablet, but purchases from unverified sources carry legal, safety and quality risks.

Telehealth services sometimes bundle consultation and medicine supply for a fixed fee. When comparing costs, include consultation, pharmacy dispensing and shipping fees, and confirm that the prescriber and supplier are operating within Australian law.

[Move back to the very top](#)

The regulatory environment for imports in Australia

- PDE5 inhibitors such as vardenafil, sildenafil, tadalafil and avanafil are Schedule 4 prescription medicines somewhere in Australia. Dapoxetine and most SSRIs are also Schedule 4.

- The fixed dose tablet sold as Super Levitra is not listed on the ARTG. Community pharmacies cannot routinely supply unapproved fixed combinations.
- You must hold a valid prescription from an Australian registered prescriber to lawfully obtain Schedule 4 medicines Australian territory.
- For unapproved medicines, prescribers can explore the TGA Special Access Scheme or Authorised Prescriber pathways, but access is not automatic and supply can be limited.
- The Personal Importation Plan allows individuals to import up to a three month supply for personal use in many cases if they have documentation such as a prescription or letter from their doctor. Customs may request evidence and rules change, so check current TGA and Australian Border Force guidance before ordering overseas.
- Buying prescription medicines from overseas websites can breach Australian law and increases the chance of counterfeit or substandard products. Advertising of prescription medicines is strictly regulated.

This summary is general information and not legal advice. For current rules consult the Therapeutic Goods Administration (TGA), Australian Border Force or a legal professional.

[Up to the page top](#)

Essential Frequently Asked Questions

How long until the medicine works and how long does the effect last?

Vardenafil is usually effective within 30 to 60 minutes and effects can last around 4 to 6 hours. Dapoxetine peaks at 1 to 3 hours and is used on demand to help with premature ejaculation.

Can I take it with food or alcohol?

It can be taken with or without food, but a heavy, high fat meal may slow onset. Keep alcohol intake low because alcohol increases the risk of dizziness and may impair sexual function. Avoid grapefruit products unless your clinician says otherwise. See [Safe Handling and Use](#) and [Cornerstone Interactions among users](#) to get more information.

Is this medicine listed on the PBS for subsidy?

Most erectile dysfunction treatments are not listed for routine use on the PBS. See [Projected Prices for Australia \(AUD\)](#) to cover common retail ranges.

Is it okay to split the tablet to take less?

Do not split or crush tablets unless a pharmacist confirms the product is scored and intended for that purpose. Many fixed dose combinations are manufactured to be swallowed whole.

What if I am taking antidepressants?

Some antidepressants interact with dapoxetine and can increase the risk of serotonin syndrome. Always review your full medication list with the prescriber. See [Vital Partnered conversational exchanges fuel collaboration and innovation](#). see the details section for more the facts.

What should I do in the case of severe chest pain or an erection lasting more than four hours?

These are potential medical emergencies. Call the emergency line or proceed to the nearest ER right away. See [Potential Reactions and When to Seek Help](#) for any questions, see the information segment in the document.

[Take me to the zenith](#)

- [Understanding Empagliflozin: Core Concepts](#)
- [7 Possible Harmful Effects of Viagra Extra Dosage You Should Know](#)
- [Mega ED Pack: where to find generic options](#)

[Additional info about Super Levitra](#)