

Low-Cost Guide to Extra Super Viagra Online - in Australia

Extra Super Viagra Side Effects and Related Body Reactions

- [Overview and constituent parts](#)
- [The mode by which it acts](#)
- [Who this product is for](#)
- [Ways to take it](#)
- [Dosage instructions](#)
- [Safety measures and safety precautions](#)
- [Drug a mutual exchange of ideas potential](#)
- [Potential systemic incidental outcomes](#)
- [Data storage management helps organize and track items. and retrieval and handling tips](#)
- [Similar medications and an alternative path choices](#)
- [Availability and value based pricing in Australia](#)
- [The statutory framework in Australia](#)
- [FAQs](#)
- [Get more information](#)

Overall view and composition

Extra Super Viagra is a combined tablet that typically pairs sildenafil with dapoxetine. Commonly quoted strengths are 100 mg of sildenafil together with 100 mg of dapoxetine in a single fixed-dose tablet. The formulation is intended for adult males.

Sildenafil addresses erectile dysfunction while dapoxetine is a short-acting selective serotonin reuptake inhibitor used to treat premature ejaculation. Note that neither component produces sexual desire; sexual stimulation is still required for an erection to occur.

Fast links: practical administration tips are under [How to accept what comes](#); safety handbooks and guidance is in [Risk reduction measures for safety](#) and [Drug intercommunication effects](#); see [Prices and availability for the Australian region](#) and [Laws governing Australia](#) for local regulatory details.

- Primary indication: erectile dysfunction support in adult men
- Active substances: sildenafil and dapoxetine
- Typical combined strength reported: 100 mg + 100 mg per tablet

How it exerts its action

Sildenafil inhibits the phosphodiesterase type 5 (PDE5) enzyme. By potentiating the nitric oxide pathway, it promotes blood vessel relaxation in penile tissue and can assist the physiological processes needed to achieve and sustain an erection during sexual stimulation.

Dapoxetine is a rapidly absorbed SSRI designed for on-demand use. It can delay ejaculation in some men by increasing serotonin availability at synapses. The rationale for a combined tablet is to address both erectile dysfunction and premature ejaculation in selected patients under medical supervision.

Reminder: neither drug increases libido; adequate sexual arousal is required for sildenafil to be effective.

Move to [Who the target audience is](#) or back to the [High level summary](#).

Who is this for

The product is indicated for adult men with erectile dysfunction who may also have premature ejaculation. It is intended to be used when sexual stimulation is expected; taking the tablet while not sexually aroused will not produce an erection.

Ways to handle it

Follow instructions provided by your prescriber or pharmacist. General advice:

- Take one tablet with water. It may be taken with or without food, but fatty meals can slow sildenafil absorption and delay onset of effect.
- Timing: many users take the tablet about one hour before sexual activity. Some people may find onset between 30 minutes and up to 4 hours, depending on individual factors.
- Frequency: do not take more than the prescribed dose. The usual maximum is one tablet in any 24-hour period unless a clinician advises otherwise.
- If you forget a planned dose, take it only if sexual activity is still intended and it is safe for you to do so. This combination is not typically for daily use unless explicitly prescribed.

For dose selection and adjustments see [Recommendations for dosing](#).

Recommendations for dosing

Always adhere to the dose recommended by a healthcare professional. Fixed-combination tablets are often marketed with 100 mg sildenafil and 100 mg dapoxetine per unit. The standard safety limit is generally one tablet per 24 hours.

If adverse reactions occur, contact your prescriber. They may switch you to single-ingredient products to allow finer dose titration or recommend an alternative treatment. Do not cut, crush, or divide tablets to modify the dose unless the manufacturer and prescriber confirm it is acceptable.

This medicine is not for children or women. People with significant liver or kidney impairment, heart disease, psychiatric illness, or who take multiple other medicines should obtain individual advice before you use it.

Continue to [Precautionary safety measures](#) and [Adverse drug negative interpersonal dialogue and exchanges between people can shape outcomes. may pose risks](#) for verify the information ahead of the start.

Caution notes and safety guidelines

Do not take this combination tablet if any of the following apply:

- Known allergy to sildenafil, dapoxetine, or excipients in the tablet
- You are using nitrates or nitric oxide donors (for example glyceryl trinitrate, isosorbide dinitrate, nitroprusside) or the guanylate cyclase stimulator riociguat

- You are already taking another PDE5 inhibitor such as sildenafil, tadalafil, vardenafil, or avanafil
- Use of recreational nitrites (poppers) including amyl or butyl nitrite
- Your clinician has advised against sexual activity because of cardiovascular risk

Discuss risks with your doctor before use if you have or have had any of the following:

- Penile structural disorders or a history of priapism, or hematological disorders such as sickle cell disease
- Eye conditions including optic neuropathy or retinitis pigmentosa; any sudden vision problems
- Unstable cardiovascular disease, recent heart attack or stroke (particularly within 6 months), angina, arrhythmia, or uncontrolled blood pressure
- Liver or renal impairment; chronic lung disorders including pulmonary veno-occlusive disease
- Seizure related neurological disorders history, bipolar disorder, severe depression, or other psychiatric illness

Dapoxetine may increase the risk of serotonin syndrome if combined with other serotonergic medicines. Avoid or use caution with other agents that affect serotonin. If any of the above conditions exist, consult a health professional before taking this medicine.

Come back toward the [top](#).

Drug a mutual exchange of ideas potential

The following medicines and substances can interact with sildenafil and/or dapoxetine and either increase risk or reduce benefit:

- Any nitrates or nitric oxide donors: can cause a severe drop in blood pressure and are contraindicated
- Alpha-blockers and other antihypertensives: combined use can lead to symptomatic hypotension
- Potent CYP3A4 inhibitors: examples include ketoconazole, itraconazole, ritonavir and some macrolide antibiotics; these may raise drug levels and treatment side effects

- CYP3A4 inducers such as rifampin or some anticonvulsants: may reduce effectiveness by lowering plasma levels
- Other serotonergic drugs: SSRIs, SNRIs, TCAs, MAO inhibitors, linezolid, lithium, tramadol and certain migraine medicines can raise the risk of serotonin syndrome when combined with dapoxetine
- Alcohol content and other central nervous system depressants: may enhance dizziness and fainting and should be limited, especially with dapoxetine
- Grapefruit or grapefruit juice: may increase sildenafil concentrations by inhibiting CYP3A4

Not all items are shown in this list. Always verify the data. with your healthcare provider or pharmacist before starting, stopping or combining medicines or supplements.

Also check out [Safety practices and hazard controls](#) and [Potential adverse outcomes](#).

Potential reactions to therapy

Many people tolerate this combination, but adverse symptoms can occur. Below are common and serious reactions to be aware of.

Common, usually short-lived effects may include:

- Migraine with facial flushing and lightheadedness, respectively.
- Dyspepsia symptoms, nausea
- Nose blockage with congestion
- Diarrhoea

Pursue urgent medical assistance for serious signs such as:

- Allergic reactions including swelling of the face, lips, tongue or throat, widespread rash or breathing difficulty
- Chest ache, loss of consciousness, and a fast or irregular heart rate.
- Sudden vision loss or significant reduction in vision; sudden hearing loss or ringing in the ears
- New weakness on one side of the body, slurred speech or other stroke-like symptoms
- Prolonged or painful erection lasting more than 4 hours (priapism)

- Severe dizziness, confusion, or seizure

If you are concerned about symptoms or they persist, contact your healthcare professional.

Navigate back to the previous page [top](#) or look over [Caution and safety measures](#).

Data temporary storage helps stage tasks before final processing. tips

- Keep tablets in their original packaging away from direct sunlight, damp locations and sources of heat
- Recommended optimized storage layouts improve overall efficiency.: around 25 C; short-term optimized storage layouts improve overall efficiency. between 15 C and 30 C is usually acceptable
- Store out of reach and sight of children and pets

For full product details see [More information can be found](#).

Other ways to treat and lookalike medicines

Depending on individual needs and medical history, clinicians may recommend different approaches. Options include:

- Tadalafil (Cialis and generics): 10 mg to 20 mg as needed, or low-dose daily regimens such as 2.5 mg to 5 mg for continuous effect. Duration can be up to 36 hours.
- Avanafil - Spedra: available in some markets at 50 mg to 200 mg; onset may be faster for some users.
- Vardenafil (Levitra and generics): typically 10 mg to 20 mg with similar onset to sildenafil.
- Sildenafil alone (Viagra and generics): available in 25 mg, 50 mg, 100 mg strengths.
- Dapoxetine alone (Priligy and generics): 30 mg or 60 mg for premature ejaculation; may be used with a PDE5 inhibitor under supervision.
- Non-drug approaches: topical local anesthetic creams or sprays for premature ejaculation, vacuum erection devices, intracavernosal injections, penile implants, lifestyle modification, and psychosexual therapy where appropriate.

In Australia, prescribers often prefer separate single-ingredient products so doses can be tailored to the patient. Fixed-dose combination tablets may be less commonly stocked.

Cross compare the two estimated price figures at [Prices and availability for the Australian region](#).

Pricing and availability across Australia

Most erectile dysfunction medicines are not subsidised by the Pharmaceutical Benefits Scheme (PBS) for ED portents, so private purchase is common. The following are approximate private-market subscription pricing ranges in Australian dollars (AUD) per tablet. Prices vary by brand, pack size and supplier.

- Generic sildenafil (50 mg-100 mg): about AUD 0.80 to 6 per tablet depending on pack size and supplier
- Brand Viagra (50 mg-100 mg): commonly AUD 12 to 35 per tablet
- Tadalafil generics (10 mg-20 mg): roughly AUD 2 to 10 per tablet; branded Cialis is usually higher
- Vardenafil generics (10 mg-20 mg): around AUD 3 to 12 per tablet where available
- Spedra as Avanafil: private prices often exceed AUD 10 per tablet
- Dapoxetine (30 mg-60 mg): roughly AUD 7 to 22 per tablet privately
- Fixed combination tablets like Extra Super Viagra: limited Australian supply; online sellers sometimes list AUD 4 to 20 per tablet before shipping and customs; legal import rules apply

Actual cost depends on pharmacy dispensing fees, telehealth consultation charges, concessions, and whether the product is imported. Sildenafil remains PBS-listed for pulmonary arterial hypertension but not usually for ED.

See [Substitute possibilities](#) or [The legal framework and standards in Australia](#) for more on supply and importation.

The legislation governing Australia

Key regulatory points to consider when obtaining these medicines in Australia:

- Prescription requirement: Sildenafil, dapoxetine and other PDE5 inhibitors are prescription-only (Schedule 4). A valid prescription from a registered prescriber is required for lawful supply.
- Dispensing: Australian pharmacies dispense prescription medicines with labelled instructions and may provide counselling.
- Online purchases within Australia: legal when supplied by a licensed Australian pharmacy that requests and verifies a valid prescription.
- Personal importation: under the Therapeutic Goods Administration (TGA) Scheme for Personal Importation you may import a limited supply for personal use with a prescription. Documentation and original packaging are important; non-compliant imports can be seized.
- Compounding: some pharmacies may compound single-ingredient products where clinically appropriate; compounded preparations still require a prescription.
- Counterfeit risk: unregulated overseas websites can sell fake or substandard products. Prefer Australian-registered pharmacies and consult with a professional for advice.
- Driving and machinery: dapoxetine can cause dizziness or fainting in some people. Avoid driving or operating machinery until you know how it affects you.

If you are unsure about legal requirements or supply options, consult your doctor or pharmacist for advice tailored to your circumstances.

Jump back to the [top](#).

Frequently queried questions

Short answers to common questions; follow the links to get more detail.

- Will it work without arousal? No. Sexual stimulation is necessary for sildenafil to produce an erection. See [How it functions in use](#).
- Should I limit alcohol while taking this medication? Limit or avoid alcohol, particularly because dapoxetine can increase dizziness and fainting risk. See [Medication potential hazards in intercommunication](#).
- How quickly does it act? Many people notice onset within 30 to 60 minutes; a high-fat meal may delay absorption. See [How to take action it calmly](#).

- What should I do about treatment side effects? Stop and seek medical attention for severe or worrying symptoms. For common reactions see [Likely temporary side effects](#).
- Is a prescription mandatory in Australia? Yes, a prescription is required. See [Legal rules in Australia](#).

More details are available

Principal active constituents: sildenafil and dapoxetine. Intended for erectile dysfunction and for delaying ejaculation in men; both require appropriate medical assessment prior to use. Physical storage management helps organize and track items. safety guidelines can be found in the guidelines. [Global flexible storage options adapt to changing needs. care guidance](#). For adverse reactions consult [Reactions you may notice](#). To review other treatment choices and indicative costs, see [Substitute medicines and analogous drugs](#) and [Current prices and current availability in Australia](#).

Return again to the [the page header](#).

- [P Force Fort - Uses, Likely Potential Adverse Reactions, and Extra Facts](#)
- [A primer on Lisinopril: what it is and FDA approval?](#)
- [Viagra Jelly Generic vs Brand Formulations: Efficacy Assessment](#)

[Additional info about Extra Super Viagra](#)