

Coupon Codes for Super Levitra: Prices and Savings - in the UK

Who is eligible to take Super Levitra and who is not

- [Summary](#)
- [Envisioned uses](#)
- [Who is not supposed to take it?](#)
- [Connections and clear the act of connecting with otherss reduce confusion.](#)
- [A guide to using](#)
- [Dose control and adjustments](#)
- [Adverse reactions](#)
- [Warehouse access to storage space determines how quickly items can be retrieved.](#)
- [Alternative solutions](#)
- [Standard price structure in the United Kingdom](#)
- [UK law and regulatory guidance notes](#)
- [For more information](#)

Summary of the material

The product commonly sold as Super Levitra combines two active molecules in one tablet: vardenafil, a phosphodiesterase type 5 inhibitor used to assist erections, and dapoxetine, a short-acting serotonin reuptake inhibitor intended to reduce premature ejaculation. This fixed-dose combination is not authorised in the United Kingdom.

Main points at a glance:

- Primary target: erectile dysfunction; the dapoxetine part aims to help with early ejaculation.
- Usual marketed strength: vardenafil 20 mg plus dapoxetine 60 mg per tablet in many online listings.
- Ingredients combined: vardenafil and dapoxetine in a single oral dose.
- Regulatory note: no UK marketing authorisation exists for a combined tablet of these two actives. Access the section that covers legal provisions at [UK compliance with laws and regulations policy framework risk guidance notes](#) before acquiring or using such a product.

Use the links to go straight to [Target uses](#), [Conversational the back and forth between peoples can be viewed as exchanges of data and details](#), or [Alternative options](#).

Envisioned uses

Manufacturers or sellers usually promote the tablet for men experiencing erectile dysfunction who also want assistance delaying ejaculation. Vardenafil addresses erectile capability while dapoxetine is an oral therapy licensed in some countries for premature ejaculation.

Because no combined product is licensed in the UK, a UK prescriber should determine whether prescribing licensed single-ingredient medicines separately would be safer and more appropriate. If you have therapy is not indicated due to these conditions or uncertainties, consult the section [Who must not take this?](#)

If combined therapy is not suitable, clinicians commonly prescribe vardenafil or tadalafil for erectile problems and, where appropriate, prescribe dapoxetine alone for premature ejaculation. Do a comparison these options under [Optional routes](#).

How to utilize this

Follow the guidance laid out by your healthcare professional. The items below are general advice and do not replace an individual consultation.

- Take a single tablet about one hour before planned sexual activity. Some people may notice effects earlier or later.
- A heavy or fatty meal may delay absorption and slow onset of effect.

- Do not take more than one tablet in 24 hours. In certain cases, for safety reasons, a prescriber may recommend spacing doses further apart, for example once every 48 or 72 hours. See [Dosage change and adjustments](#).
- Avoid grapefruit or grapefruit juice unless your clinician permits it, because grapefruit can raise drug concentrations. See [Connections and interactions among individualss create opportunities for collaboration.](#)
- If you forget a planned dose and sexual activity is still anticipated, take the medicine when you remember provided the 24 hour rule is observed; never double the dose to catch up.

To secure this, see for safe keeping [Archival modular storage systems allow easy reconfiguration.](#). If any exclusion criteria may apply to you, read [Who should not be allowed to take this?](#).

Dosing plans and adjustments

Only a prescriber can tailor dose and dosing interval. Factors that commonly require dose modification include age, liver or kidney function, cardiovascular status and interacting medicines.

- Older patients (for example those over 65) and people with moderate to severe liver impairment often require lower doses and longer intervals between doses.
- Concurrent use of alpha-blockers for benign prostatic hyperplasia or blood pressure may increase the risk of low blood pressure; clinicians may start at a reduced dose and monitor closely.
- Co-administration with drugs that raise vardenafil or dapoxetine exposure (for example strong CYP3A4 inhibitors) may lead a prescriber to limit dosing frequency to once every 72 hours or to select an a substitute choice.

If you are unsure about the correct dose for you, contact your prescriber. Verify the data absolute exclusions details [Who should not be taking this?](#).

How to preserve items in temporary storage helps stage tasks before final processing.

- Store at room temperature around 25 C (77 F). Short deviations from this are usually acceptable as stated on the packaging.
- Keep this product out of reach and out of sight of children.

- Protect the tablets from moisture and excessive heat. Leave them in their original blister until used.
- Do not use the tablets after the expiry date printed on the pack.

Return to the point you were at [How to make use of](#) or go back to the [First look at the apexic](#).

Who should not take this medicine

Do not take this combined tablet if any of the following apply. If in doubt, seek medical advice immediately.

- You are using nitrates such as glyceryl trinitrate or isosorbide, or you use recreational nitrites (poppers). Combining nitrates with a PDE5 inhibitor can cause a dangerous fall in blood pressure.
- A clinician has advised you to avoid sexual activity because of cardiac risk.
- You have unstable or severe cardiac disease, recent heart attack or stroke within the past 6 months, severe heart failure, or life-threatening arrhythmias.
- You have severe liver impairment or are on dialysis for severe renal failure.
- You have very low blood pressure or uncontrolled high blood pressure.
- You are allergic to vardenafil, dapoxetine, or any tablet ingredient.
- You have inherited retinal disorders such as retinitis pigmentosa.
- You have a congenital or acquired long QT interval syndrome or are taking medicines known to prolong the QT interval significantly.
- You are already using a different PDE5 inhibitor (for example sildenafil, tadalafil, avanafil, or single-ingredient vardenafil).
- You are taking monoamine oxidase inhibitors (MAOIs), have used MAOIs in the last 14 days, or you take thioridazine; dapoxetine is contraindicated with these agents.

Tell the prescriber about any history of blood disorders, penile curvature or deformity, priapism, eye disease, hearing problems, peptic ulcer, seizures, pulmonary hypertension, syncope, bleeding disorders or a family history of long QT syndrome.

For the reciprocal contact between individuals among teammates fuel progress. with medicines and food consult [Social interplay between parties between people can shape outcomes.](#) If any exclusion above applies to you, do not take this medicine and consult a physician.

Food can influence medicines.

The combination of vardenafil and dapoxetine can interact with numerous drugs and some foods, increasing indirect effects or reducing effectiveness.

- Nitrates and nitroprusside: absolutely contraindicated because of the risk of severe hypotension. See [Who should not be given it?](#).
- Alpha-blockers such as doxazosin or tamsulosin: may cause symptomatic low blood pressure; start low and monitor.
- Strong CYP3A4 inhibitors (for example ketoconazole, itraconazole, ritonavir, some macrolides): can raise levels of either active and increase adverse reactions.
- Strong enzyme inducers like rifampicin: may lower vardenafil levels and reduce its effect.
- Certain antiarrhythmics (for example amiodarone, sotalol) and other QT-prolonging agents: potential for cardiac rhythm problems.
- Other serotonergic drugs (for example SSRIs, SNRIs, triptans, tramadol, linezolid, St Johns wort flower, lithium): combination with dapoxetine can raise the risk of serotonin syndrome; MAOIs and thioridazine must be avoided. See [Who ought not to take this?](#).
- Grapefruit and grapefruit juice: may increase systemic drug levels; avoid unless your clinician says it is safe. See [Best practices for using.](#)
- Ethanol: may worsen dizziness, fainting and impair judgment, particularly when combined with dapoxetine.

This list is only a portion of the total. Always provide your clinician with a full list of prescribed medicines, over-the-counter remedies, herbal preparations and supplements.

Possible unwanted effects

Many users tolerate treatment well, but undesired reactions can occur. If you have concerns, schedule an appointment with a healthcare professional.

Frequent drug adverse effects, typically mild:

- A throbbing a mild headache and facial flushing
- The symptom of dizziness or lightheadedness
- Congestion in the nose or a runny nose
- Stomach discomfort including indigestion or nausea
- Lack of saliva or difficulty sleeping; diarrhoea is more often linked to dapoxetine

If you notice any alarming signs, seek urgent medical assistance any of the following:

- Signs of a severe allergic reaction such as widespread rash, swelling of the face, lips or throat, or breathing problems
- Chest pain, fainting, or a very fast, slow or irregular heartbeat
- A sudden loss of vision in one eye or in both eyes., or sudden hearing loss with or without ringing in the ears
- An erection lasting longer than four hours (priapism) or a painful erection
- New seizures, sudden severe head discomfort or vomiting, numbness or severe weakness, or sudden confusion or memory loss

For regulatory obligations concerning adverse reactions in the UK consult the [UK law and regulatory briefing notes](#) part of the document. Return to [Who should not be given it?](#) for exclusion criteria.

Substitute therapies and comparable drugs

If a combined unlicensed tablet is not appropriate, the following licensed options may be considered after clinical review.

- Tadalafil: a longer acting PDE5 inhibitor with onset in about 30 to 60 minutes and effect lasting up to 36 hours; daily low dose options are available.
- Vardenafil as a single ingredient: onset typically 30 to 60 minutes with a duration around 4 to 6 hours.
- Sildenafil: well established PDE5 inhibitor; common onset 30 to 60 minutes and effect for around 4 hours. In the UK a 50 mg pharmacy-only supply (Viagra Connect) can be provided after pharmacist assessment.

- Avanafil: a newer PDE5 inhibitor reported to act in as little as 15 to 30 minutes for some patients, lasting up to about 6 hours.
- Dapoxetine on its own: licensed in some countries for premature ejaculation and usually dosed at 30 or 60 mg as needed subject to suitability checks.
- Alprostadil: local treatments such as intraurethral or intracavernosal options for men who cannot use or do not respond to PDE5 inhibitors.
- Penile negative-pressure erection devices: a non-drug mechanical option that can be used alone or in combination with other therapies.
- Non-pharmacological approaches: lifestyle changes, weight loss, smoking cessation, reduced alcohol intake and psychosexual counselling can all improve sexual function.

For indicative private prices in GBP see [Regular UK range of costs](#). Always discuss the options with a prescriber to identify the safest and most effective choice.

Common price bands in Britain (private, per tablet)

Prices depend on brand, strength, pack size and whether you buy privately online or from a high street pharmacy. The NHS provides restricted access to ED medicines, so many people pay privately.

- Generic sildenafil 50-100 mg: roughly 1 to 3 pounds per tablet; branded Viagra typically 8 to 10 pounds.
- Generic vardenafil 10-20 mg: roughly 2 to 6 pounds per tablet.
- Generic tadalafil 10-20 mg: about 1.50 to 3 pounds per tablet; daily 5 mg may cost around 0.80 to 2 pounds each depending on pack size.
- Avanafil: Spedra 100-200 mg: often around 6 to 9 pounds per tablet.
- Dapoxetine 30-60 mg (licensed single-ingredient product): commonly 2 to 5 pounds per tablet.
- Unlicensed combination products marketed as Super Levitra: prices vary widely online, for example 3 to 10 pounds per tablet; quality, authenticity and legality are uncertain.

Additional fees such as online consultation or private prescription charges commonly add 10 to 30 pounds per order. Devices like vacuum pumps range from about 30 to 120 pounds depending on model.

Weigh up the options at [Comparable drugs and nontraditional medical treatments](#) and examine the data the legal position in [UK statutes and notes on regulations](#).

UK legislation and notes on regulatory changes

- Prescription status: Vardenafil and dapoxetine are prescription-only medicines (POM) in the UK. A UK-registered prescriber must assess suitability and provide a prescription for legal supply.
- No licensed combination: There is no UK marketing authorisation for a single tablet that contains both vardenafil and dapoxetine. Products offered online claiming this formulation are likely unlicensed.
- Risks of import: Unlicensed imports may be detained by Border Force and can pose safety risks. The Medicines and Healthcare products Regulatory Agency (MHRA) warns about falsified or unsafe medicines bought online.
- Buying online: Use pharmacies registered with the General Pharmaceutical Council (GPhC) and corroborate the evidence for clear contact details. Legitimate UK services will show GPhC registration the available information.
- Pharmacy supply: Some products, for example sildenafil 50 mg as Viagra Connect, can be supplied after pharmacist assessment without a prescription. Other PDE5 inhibitors remain prescription-only.
- NHS prescribing: NHS access to ED medicines is restricted and varies by local policy; many patients obtain treatment through private consultations.
- Age and assessment: You must be at least 18 and complete a health assessment so a prescriber can check for red flags against use of this medication such as nitrate use.
- Driving and machinery: If you experience dizziness, fainting or visual disturbance, do not drive or operate machines until these effects have cleared.

For safety concerns and rules affecting supply see [Who should not take it?](#) and [Human people engage through a series of the dynamics of interactions.](#). Return to the position you left [Macro view](#).

More details

Bioactive ingredients summary: vardenafil enhances erectile response by inhibiting PDE5, increasing penile blood flow during sexual stimulation. Dapoxetine is designed to delay ejaculation by potentiating serotonergic activity in the central nervous system.

If you have specific questions about suitability, dosing, or potential interplay between parties create context for decisions., speak to your doctor or pharmacist. In case of severe or worrying symptoms seek emergency care.

Read more on [follow-on effects](#), return to [how to use this product](#) or balance with in a comparison [secondary treatment options](#).

[Go to top](#)

- [Understanding Tadalafil Strong: definition and FDA approval?](#)
- [Uses, Unwanted reactions, and More for Cialis Black](#)
- [Viagra Extra Dosage: Secret Scripts for Patients from Physicians](#)

[Additional info about Super Levitra](#)

© 2026 Kimbrell Veterinary Clinic PC - [Home page](#) - [We're on the map](#)