

Quick Guide to Extra Super Viagra: Uses, Dosing, Price, and Safety - in the UK

Is Extra Super Viagra a viable option for you?

- Comprehensive the essentials at a glance and composition
- The target audience for this
- How to take responsibility
- Strength and dosing details
- Important alerts for safety
- Collaborative the interactional flow among participantss among teammates fuel progress. to validate and verify
- Potential adverse outcomes
- Archival cold storage is used for long term preservation of rare items. tips
- Cost guide and other options (GBP)
- What is the legal recognition status in the United Kingdom? and access
- More details

A concise a short outline of the key ideas and its composition

Product name used in some markets: Extra Super Viagra. It is presented as a single tablet combining two active agents: sildenafil and dapoxetine.

Sildenafil acts as a phosphodiesterase type 5 inhibitor to help with erectile function. Dapoxetine is a short-acting selective serotonin reuptake inhibitor normally prescribed for premature ejaculation in certain countries.

Reported formulations commonly promoted online claim 100 mg sildenafil plus 100 mg dapoxetine per tablet, though availability, exact content, and quality can vary widely by supplier and are not guaranteed.

Fast jump to navigation [keep in mind the safety safety advisories](#), see known [medication the risks of the give and take of interaction](#), or view [an alternate path medical therapies and approximate charge estimates](#). For information about the law, see [United Kingdom status of lawful authorization and how to access](#).

People this feature is aimed at

This combination is promoted for adult men who have erectile dysfunction and who also report problems with early ejaculation. Note that sexual stimulation is still required for sildenafil to work.

In the United Kingdom, preparations that combine sildenafil and dapoxetine in a fixed dose are not authorised. Dapoxetine itself is available only on prescription when clinically indicated for premature ejaculation.

[Navigate back to the an overarching view of the material page](#)

How to take steps this product

Follow the instructions from a qualified prescriber. General practical points are listed below.

- Timing: Most people take sildenafil-containing tablets about one hour before sexual activity; some can take them from 30 minutes to up to four hours beforehand depending on the product and individual response.
- Food: You may take the tablet with or without food. A heavy, high-fat meal can delay how quickly it starts to work.
- Frequency: Do not take more than one dose in any 24-hour period unless a prescriber advises otherwise.
- Dose not taken today before planned activity: If you forget to take it before sexual activity, take the dose when you remember but do not double up to make up for a dose was omitted.
- Liquor: Avoid excess alcohol because it can reduce erectile response and increase dizziness; alcohol combined with dapoxetine may raise the chance of fainting or other detrimental health effects.

For guidance on strengths and adjustments, see [Dosage and strengths](#).

Dosing amount and strength

Only take medicines at the dose prescribed by your healthcare professional. Many online sellers list tablets containing 100 mg sildenafil and 100 mg dapoxetine, but this is not an approved combination in the UK.

Licensed single-ingredient dosing in the UK (for comparison):

- Sildenafil (single-ingredient products): typical on-demand dose is 50 mg taken approximately 1 hour before sex. Usual range is 25 mg to 100 mg depending on effect and tolerability.
- Dapoxetine (single-ingredient products): the usual starting dose is 30 mg, which may be increased to 60 mg if needed and tolerated. A single 100 mg dose of dapoxetine would exceed licensed UK doses.

Do not alter tablet doses, crush or split tablets you are not instructed to split, or combine this product with other ED or SSRI medicines without medical supervision. If you feel the dose is too strong or not effective, discuss options in dialog with your prescriber.

Additional reading [Caution and safety notices](#) and [Social exchanges](#).

Important remember the safety warning notices

Do not use this product if

- You are taking nitrates in any form, for example nitroglycerin, isosorbide mononitrate, or sodium nitroprusside.
- You inhale or inject recreational nitrites often called poppers (such as amyl nitrite or butyl nitrite).
- You are already using another PDE5 inhibitor like tadalafil or vardenafil, or any product that contains sildenafil.
- A healthcare professional has told you to avoid sexual activity because of cardiac risk.
- You know you are hypersensitive to sildenafil, dapoxetine, or any ingredient in the tablet.

Tell your doctor or pharmacist before using if you have

- Any prescription or over-the-counter medicines, herbal remedies, or supplements.

- A history of liver or kidney disease, uncontrolled blood pressure, stomach ulcers, bleeding disorders, heart disease including recent heart attack or stroke within the last six months, or serious arrhythmias.
- Blood disorders such as sickle cell disease, leukemia, or multiple myeloma, or a tendency to prolonged erections.
- Penile structural problems such as Peyronie disease or previous prolonged erections.
- Eye conditions including optic neuropathy or retinitis pigmentosa, or sudden vision loss in the past.
- A history of psychiatric illness, mania, bipolar disorder, recurrent depression, fainting spells, or seizures; these are particularly relevant because of dapoxetine.

Please consult a healthcare professional before you begin or cease any medication. See [Engagements and exchanges that require checking](#) for drugs and foods to avoid.

Double check of communication in actions in the user journey shape satisfaction.

Several medicines and substances can increase unfavorable effects or change how well either component works. Always disclose everything you take to your prescriber.

- Strong CYP3A4 inhibitors such as ketoconazole, itraconazole, certain macrolide antibiotics (for example erythromycin and clarithromycin), protease inhibitors, and grapefruit juice can raise blood levels of sildenafil and dapoxetine.
- Alpha-blockers (for example doxazosin or tamsulosin), other antihypertensives, or nitrates can cause dangerous blood pressure falls if combined with sildenafil.
- Enzyme inducers like rifampicin, carbamazepine, phenytoin, or bosentan may lower concentrations and reduce effectiveness.
- Other serotonergic drugs including SSRIs, SNRIs, MAO inhibitors, linezolid, lithium, tramadol, triptans, or The medicinal herb St Johns wort can increase the risk of serotonin syndrome when used with dapoxetine. Dapoxetine must not be taken with thioridazine or within recommended washout periods of MAOIs.
- Recreational drugs and heavy alcohol use increase the risk of dizziness, fainting, and impaired judgment.

This is not a full inventory. Contact a clinician or pharmacist before changing any medicine. Return to the previous page [Environmental warnings on safety](#).

Possible unintended effects

Many people tolerate treatment well but adverse reactions can occur. Below are common and serious effects to be aware of.

Common, mild in most scenarios

- Head pressure and a red face
- Nausea, indigestion, or heartburn
- Nasal blockage or a runny nose
- The runs or dry mouth
- Dizzy spell

Seek immediate medical attention if you experience

- Anaphylactic or severe allergic reaction: widespread rash, swelling of face or throat, breathing difficulty
- Chest pain, collapse, fainting, or a very fast or irregular heartbeat
- Sudden partial or complete loss of vision in one or both eyes
- Sudden decrease or loss of hearing, often with ringing in the ears
- Seizure, prolonged severe dizziness, or new confusion
- Painful erection lasting more than four hours (priapism) — this is an emergency
- Signs of stroke such as sudden weakness or numbness on one side of the body, slurred speech, or drooping of the face

Should any symptom cause concern, consult your healthcare provider without delay. See also related links [Cooperative exchanges to check up on](#).

Remote modular storage systems allow easy reconfiguration. recommendations

- Keep tablets in their original packaging, protected from moisture and light. Avoid storing in a bathroom where humidity is high.

- Store at around 25 C (77 F); short variations are usually acceptable but avoid prolonged high temperatures.
- Keep this item out of reach and out of sight for kids and animals.

Please refer back to the earlier content [Safety advisories](#) for more comprehensive health and safety measures.

Choices and value based pricing guide in the UK (GBP)

Licensed single-agent therapies are the usual clinical route in the UK. Prices depend on whether a product is supplied on the NHS, privately by prescription, or bought as a pharmacist-supplied product.

- Sildenafil (generic or Viagra): generic options on private prescription can be approximately 0.20 to 1.50 GBP per tablet for 50 to 100 mg strengths. Viagra Connect 50 mg, available after a pharmacist consultation, is often around 15 to 25 GBP for a pack of 4 tablets.
- Tadalafil (Cialis or generic): private generic tablets of 10 to 20 mg commonly range from about 0.60 to 2.00 GBP each; brand prices are higher.
- Vardenafil (Levitra or generic): private prices typically fall in the region of 1.50 to 4.00 GBP per 10 to 20 mg tablet.
- Avanafil; Spedra: often supplied as brand-only and priced roughly 4.00 to 10.00 GBP per tablet depending on strength.
- Dapoxetine (Priligy, generic): for premature ejaculation on private prescription, expect around 10.00 to 25.00 GBP per 30 to 60 mg tablet depending on supplier and quantity.
- Fixed-dose combinations (sildenafil plus dapoxetine), including products promoted as Extra Super Viagra, are not licensed in the UK; some overseas sellers advertise prices around 2.00 to 8.00 GBP per tablet, but quality and legality are uncertain and such products are not recommended.

Other PDE5 inhibitors occasionally discussed internationally include udenafil and lodenafil, but these are not commonly used or licensed in the UK. For safe and legal supply, consult a UK-registered prescriber or pharmacist.

Identify the better choice options and see legal details at [UK legal framework and access](#).

Britains legal standing and protected access

- Sildenafil is normally a prescription-only medicine in the UK. A specific 50 mg sildenafil product, marketed as Viagra Connect, is available as a pharmacy medicine (no prescription) after a pharmacist assessment.
- Dapoxetine is a prescription-only medicine in the UK.
- Fixed-dose combinations that pair sildenafil and dapoxetine are not authorised for sale or supply in the UK. Healthcare professionals typically prescribe licensed single-ingredient products where clinically appropriate.
- Importation and supply of unlicensed medicines are controlled by the MHRA. Personal importation can lead to seizure or other enforcement, and it is generally unlawful for UK businesses to sell unlicensed combination products to the public.
- These medicines are intended for adult use only; they are not for use in children or by women unless a prescriber advises otherwise. Do not drive or use tools or machinery if you become dizzy or have visual disturbances.

For safe and lawful access, use UK-registered prescribers and drug retail outlets. If in doubt, ask a pharmacist or GP for advice on licensed ancillary choices and proper surveillance of system activity.

[See available substitute options coupled with their price tags.](#) or proceed to [a survey of the main points.](#)

Supplementary details

Active subassemblies discussed here are sildenafil and dapoxetine. Use medicines exactly as directed by a healthcare professional and read any patient information leaflet provided with the product.

If you experience any of the warning signs described under [Possible effects on your body](#) or have concerns about clear mutual interactions reduce confusion. listed in [Interactions among users to validate and verify](#) contact medical professionals promptly.

Handy quick links: [Safety](#) | [Connections along with the quality of interacting as a shared activity matters for trust.](#) | [Dosage](#) | [Status of legality](#)

[Return to the location page top](#)

- [Modalert different dose levels: forms, strength levels, and how to take notes efficiently](#)
- [Is Lisinopril the generic formulation as efficacious as the branded one?](#)

- [Propecia: dosage forms, strength variants, and intake instructions](#)

[Additional info about Extra Super Viagra](#)

© 2026 Kimbrell Veterinary Clinic PC - [Home page](#) - [We're on the map](#)