

Online portal: Super Cialis at the lowest US price

Super Cialis: eligibility and contraindications

- [Intent and a top line summary](#)
- [What happens and how the workflow comes together](#)
- [Ways to use](#)
- [Dosing directives merged with the dose was skipped](#)
- [Archive storage capacity determines how much you can keep. technique recommendations](#)
- [Who should not be permitted to use it](#)
- [Information you should provide to your clinician.](#)
- [Medications and substances clear interplay between parties reduce confusion.](#)
- [Critical safety engineering considerations](#)
- [Possible collateral effects](#)
- [Alternative drugs and similar pharmaceutical options](#)
- [US price scale \(approximate\) \(USD\)](#)
- [A legal status and purchasing in the United States](#)
- [Look at the additional notes below](#)

Executive view and rationale

Product name: Super Cialis

The active ingredients: tadalafil (a phosphodiesterase type 5 inhibitor) combined with dapoxetine (a short-acting selective serotonin reuptake agent).

Intended use: Designed for adult men who require help achieving an erection and who also seek on-demand treatment for premature ejaculation.

Common formulation: One marketed combination contains about 20 mg tadalafil together with 60 mg dapoxetine in a single tablet, although formulations may vary by manufacturer.

Expected duration: Tadalafil can support erections for many users for up to 36 hours when sexual stimulation occurs; dapoxetine works on ejaculatory latency for on-demand use.

Note: Dapoxetine is not approved by the U.S. Food and Drug Administration for use in the United States. The section addressing legal concerns can be found at [Legality and purchases in the United States](#) for more information, see the policy page.

Quick access links: more options are described at [Substitute therapies and comparable drugs](#), and price point guidance is at [US quote pricing in ballpark terms](#).

[Take me to the number one spot](#)

How the system works

Tadalafil component: inhibits the PDE5 enzyme in penile tissue which promotes relaxation of smooth muscle and increased blood flow during sexual arousal, supporting erection formation.

Dapoxetine component: briefly increases serotonergic activity in the central nervous system to help delay ejaculation when taken as directed in eligible men.

Important: The erectile effect requires sexual stimulation to be effective. For timing and administration details, consult [Ways to use this](#) and [Dosing and forgot to take my doses](#).

[Up to the capstone](#)

A quick guide to using

Always follow the instructions provided by your prescribing clinician. Individual medical factors may change the recommended approach.

- Swallow each tablet with a full glass of water. Avoid grapefruit and grapefruit juice because they can alter blood levels of tadalafil.
- It may be taken on an empty stomach or after a light meal; heavy meals may delay onset.

- On average, tadalafil begins to act within about 20 minutes for some users, while dapoxetine may require roughly 40 to 60 minutes to reach effect.
- Most guidance suggests taking the dose around 30 to 60 minutes before planned sexual activity, but follow your prescriber's timing advice.
- Limit alcohol; combining alcohol with this treatment can increase risk of dizziness, fainting, and other potential side effects.

[Move to top](#)

Dose timing and forgot to take the dose

Take the dose your prescriber selected. Do not use more than the prescribed amount or more frequently than recommended for each sexual encounter.

If you miss a planned dose and still intend to have sexual activity, take the tablet when you remember if timing still allows. If it is too close to the next planned use, skip the missed one; do not double up.

Patients with kidney or liver impairment, or those on interacting medicines, may need dose adjustments. Schedule a consultation your healthcare provider for personalized dosing. See [Items you should mention to your clinician](#) for details that may affect dose selection.

[Return for a second look top](#)

Global containers and shelves organize storage space effectively. tips and recommendations

Keep in a cool, dry place at normal room temperature (about 68 to 77 F or 20 to 25 C). Protect tablets from strong light, moisture, and heat. Store in the original container and away from children and pets.

Dispose of tablets that are past their expiry or no longer required in keeping with local regulations or take-back programs at pharmaceutical retailers.

[Go back to top](#)

Who should not make use of it

- Anyone concurrently using nitrates (for chest pain) or organic nitrites (recreational "poppers") due to life-threatening blood pressure drops.

- People with known hypersensitivity to tadalafil, dapoxetine, or any tablet ingredient.
- Those with severe hepatic impairment or end-stage renal disease without specialist advice.
- Individuals with recent serious cardiovascular events such as myocardial infarction within the past 90 days, recent stroke, or unstable angina.

If any of these apply to you, do not start the medicine without urgent medical review.

[See conditions to discuss with your prescriber](#) to give further context.

[Return to the starting point page top](#)

Signs to report to your clinician.

- All prescription, over-the-counter, and herbal medicines you take.
- Any existing heart disease, angina, heart rhythm problems, or history of heart attack or stroke.
- Ailments of the liver or kidneys, bleeding disorders, peptic ulcer disease, or any blood dyscrasia such as sickle cell disease.
- A history of severe allergic reactions to medicines or foods.

Inform your prescriber about fertility plans, use of recreational drugs, and any psychiatric medicines, including SSRIs or other antidepressants.

[Go back to the crest of the page](#)

Pharmacologic the process of people engagements across departments improve alignment. between drugs and substances

Certain combinations can cause serious problems such as large drops in blood pressure, fainting, or other adverse outcomes. Discuss all medications with your healthcare provider ensure setup before use.

- Nitrates and nitroprusside are strictly contraindicated with PDE5 inhibitors due to severe hypotension risk.
- Alpha-blockers (used for prostate enlargement or high blood pressure) can add to low blood pressure and dizziness when combined with tadalafil.

- Some antifungals, macrolide antibiotics, protease inhibitors, and certain anticonvulsants can alter blood concentrations of either active ingredient and require dose review.

Combining multiple erectile dysfunction treatments or adding recreational substances can be dangerous. See [Top safety pointers](#) for more on risk mitigation.

[Return to the point you were at top](#)

Vital safety guidelines

- Common effects include lightheadedness, drowsiness, blurred vision, and dizziness. Until you know how you react, avoid driving or using heavy machinery.
- Heat, intense physical exertion, fever, or alcohol use may increase the chance of feeling faint or dizzy.
- Heavy alcohol consumption (for example, five or more standard drinks) can raise the likelihood of experiencing a headache, palpitations, and low blood pressure.
- Stop sexual activity and seek urgent medical care if you experience chest pain, sudden weakness, sudden severe pain in the head, trouble speaking, vision loss, fainting, or prolonged erection lasting more than four hours.
- This medicine does not protect against sexually transmitted infections or prevent pregnancy; use barrier protection and contraception as appropriate.
- Not intended for women, adolescents, or children under 18 years of age.

[Go to the very top](#)

Possible collateral effects

Most users experience no serious problems, and many have only mild, transient symptoms.

- Head pain and facial flushing.
- Runny nose or stuffed nose.
- Back or muscle ache in some cases.
- Dizzy spell, sleepiness, or visual disturbances.

Rare but urgent issues include priapism (a painful erection lasting more than four hours), severe allergic reactions, or breathing difficulties. If you encounter any of these, seek emergency care immediately.

Interaction-driven detrimental outcomes may occur; check [Drug and substance the act of interaction considerations](#) for agents that raise risk.

[Back to the crown](#)

Other medicines alongside their analogs

If a tadalafil-dapoxetine combination is unsuitable, prescribers can suggest different single agents or strategies tailored to needs and health status.

- Sildenafil (Viagra) for erectile dysfunction: typically effective for 4 to 6 hours; widely available in generic form at lower cost.
- Tadalafil (Cialis) alone: long duration up to 36 hours; available as daily low-dose or as on-demand dosing.
- Avanafil (Stendra): can have a faster onset for some users; duration often around 6 hours.
- Vardenafil (Levitra): similar profile to sildenafil; oral dispersible preparations exist.
- Dapoxetine (Priligy) alone: on-demand SSRI for premature ejaculation in countries where it is approved; not FDA approved in the USA.
- Topical anesthetic approaches: lidocaine or prilocaine sprays and creams to reduce penile sensitivity for PE management.
- Behavioral therapies: counseling, pelvic floor exercises, and behavioral techniques can help with premature ejaculation and sometimes reduce medication need.

Compare the two the price points for common other options in the [Approximate value based pricing for the US market](#) next section. Discuss benefits and acute side effects with your prescriber to select the best option.

[Go to top](#)

Approximate price points in the United States (USD)

Out-of-pocket costs will vary by pharmacy, quantity, brand versus generic, and insurance coverage. The ranges below are approximate cash prices per tablet or per typical single-use dose.

- Generic sildenafil (typical 50 to 100 mg): roughly \$0.50 to \$3 per dose depending on source.
- Generic tadalafil (10 to 20 mg): about \$1 to \$5 per tablet at many U.S. pharmacy outlets.
- Brand Cialis (tadalafil): can cost between \$12 and \$70 per tablet without discounts.
- Avanafil (Stendra): commonly priced around \$10 to \$30 per tablet.
- Vardenafil generics: roughly \$3 to \$10 per tablet.
- Dapoxetine: not commercially available through FDA-approved U.S. channels; international online listings often show varied rate card and uncertain quality.
- Combined products marketed as tadalafil plus dapoxetine: not FDA approved in the USA; online international prices sometimes listed at \$2 to \$8 per tablet but legality and authenticity are uncertain.

Saving tips: manufacturer coupons, pharmacy discount cards, generic substitution, and mail-order programs can substantially lower the price of FDA-approved medications obtained from licensed U.S. pharmacist shops.

[Return to the previous page top](#)

How current legal status influences purchasing in the United States

- Tadalafil is FDA approved for erectile dysfunction and requires a valid prescription from a U.S. prescriber with an active prescribing license.
- Dapoxetine is not approved by the FDA; fixed-dose combination products that include dapoxetine are not approved in the United States.
- Purchasing prescription medications legally in the U.S. requires a prescription. Telehealth services may issue prescriptions when clinically appropriate.
- Importing medicines that are not FDA approved is generally prohibited. The FDA may exercise enforcement discretion in limited personal-use scenarios, but shipments can be detained or refused and product authenticity cannot be guaranteed.

- Use only reputable U.S.-licensed medicine dispensaries. Look for accreditation such as NABP .pharmacy or Verified Internet Pharmacy Practice Sites (VIPPS). Avoid sites that sell prescription drugs without requiring a prescription.
- Possession or use of prescription drugs without a valid U.S. prescription may violate federal and state laws. When traveling, keep medicines in original labeled packaging with prescriber the facts.
- Age restrictions: erectile dysfunction medications are intended for use by adults 18 or older; minors and women should not use these products without specific medical advice.

Please note that this information is educational, not legal counsel. For case-specific guidance, consult a licensed attorney, pharmacist, or your prescriber.

[Head back to page top](#)

More notes

Signs summary: intended for adult men with erectile dysfunction; the dapoxetine component aims to treat premature ejaculation in eligible patients in jurisdictions where it is approved.

Planning and timing: allow sufficient time before sexual activity to permit both agents to reach therapeutic levels. Sexual stimulation remains necessary for the tadalafil portion to produce an erection.

Medical supervision: regular review of cardiovascular health, liver and kidney function, and concurrent medicines is recommended while using ED therapies and drugs that affect serotonin.

[Go to top](#)

- [Viagra Extra Dosage: Can it help you?](#)
- [Viagra Professional Therapy additional effects: What Else It Affects in the Body](#)
- [A primer on Cialis Sublingual: what it is and FDA approval?](#)

[Additional info about Super Cialis](#)